

Germline *SDHB* Mutations and Familial Renal Cell Carcinoma

Christopher Ricketts, Emma R. Woodward, Pip Killick, Mark R. Morris, Dewi Astuti, Farida Latif, Eamonn R. Maher

Familial renal cell carcinoma (RCC) is a heterogeneous disorder that is most commonly caused by germline mutations in the *VHL*, *MET*, and *FLCN* genes or by constitutional chromosome 3 translocations. However, for many patients with familial RCC, the genetic basis of the disease is undefined. We investigated whether germline mutations in fumarate hydratase (*FH*) or succinate dehydrogenase subunit genes (*SDHB*, *SDHC*, *SDHD*) were associated with RCC susceptibility in 68 patients with no clinical evidence of an RCC susceptibility syndrome. No mutations in *FH*, *SDHC*, or *SDHD* were identified in probands, but 3 of the 68 (4.4%) probands had a germline *SDHB* mutation. Patients with a germline *SDHB* mutation presented with familial RCC (n = 1) or bilateral RCC (n = 2) and no personal or family history of pheochromocytoma or head and neck paraganglioma. Age at diagnosis of RCC in *SDHB* mutation carriers ranged from 24 to 73 years. These findings 1) demonstrate that patients with suspected inherited RCC should be examined for germline *SDHB* mutations, 2) suggest that all identified *SDHB* mutation carriers should be offered surveillance for RCC, and 3) provide a further link between familial RCC and activation of hypoxic-gene response pathways.

J Natl Cancer Inst 2008;100:1260–1262

Renal cell carcinomas (RCC) account for approximately 2% of all cancers in the West, and they are histologically heterogeneous, with most (~80%) classified as clear cell, that is, conventional RCC (cRCC). Among the non-clear cell types, papillary (chromophilic) and chromophobic tumors are the most frequent (1). Although only 2–3% of all cases of RCC are familial, the identification of familial cases is important so that surveillance can be offered to patients and at-risk relatives and morbidity and mortality reduced (2,3). In addition, information on the molecular basis of familial RCC (eg, in von Hippel–Lindau disease) has provided the scientific basis for the introduction of novel therapeutic strategies for sporadic RCC (4).

Familial RCC may be caused by mutations in the *VHL*, *MET*, or *FLCN* genes (mutation of *FLCN* causes Birt–Hogg–Dube syndrome) or by constitutional translocations (particularly involving chromosome 3) (5–7). However, in many cases, the genetic basis of familial RCC is unknown (8,9). Germline mutations in *FH*, the gene encoding fumarate hydratase, an enzyme in the Krebs cycle, causes a hereditary leiomyoma-

tosis syndrome that is characterized by cutaneous leiomyomas and, in women, multiple early onset uterine leiomyomas (fibroids). In men and women with these mutations, type 2 papillary RCC may also occur (10,11). Germline mutations affecting another Krebs cycle enzyme, succinate dehydrogenase (SDH), can also cause an inherited tumor susceptibility syndrome. Thus, inherited mutations in *SDHB*, *SDHC*, and *SDHD* genes (encoding subunits of SDH) are associated with a high risk of head and neck paragangliomas and/or adrenal and extra-adrenal pheochromocytoma (12–15). Very rarely, germline mutations in *SDHB* have been found in patients with RCC and a personal or family history of pheochromocytoma and/or paraganglioma (16). Similarly, germline mutations in *SDHB*, *SDHC*, and *SDHD* have been detected in patients with a combination of gastrointestinal stromal tumors and paraganglioma (17).

We investigated whether 68 patients with features of nonsyndromic inherited RCC susceptibility might harbor mutations in *FH*, *SDHB*, *SDHC*, or *SDHD*. Twenty-nine of the probands had familial RCC, and 39 were without demonstrable family his-

tory of RCC (21 of the latter group had early multicentric RCC and 18 had early-onset [age at diagnosis <40 years] disease). All patients studied had 1) no clinical evidence of an RCC susceptibility syndrome and 2) normal cytogenetic analysis and normal *VHL* and *FLCN* mutation analysis. All subjects gave consent for genetic studies, and the investigations were approved by the South Birmingham Research Ethics committee. To detect mutations, the *FH*, *SDHB*, *SDHC*, and *SDHD* genes were sequenced as described previously (18). No mutations in *FH*, *SDHC*, or *SDHD* were detected. However, 3 of the 68 (4.4%, upper limit of calculated population proportion 95% confidence interval = 9.29%) probands (3.4% of probands with familial disease and 9.5% of those with multicentric RCC) had a germline *SDHB* mutation.

The mutations in these three patients were as follows. Case subject 1: A germline heterozygous nonsense mutation, p.Arg46Stop (c.136C>T, Figure 1) in exon 2 was detected in a 58-year-old male proband who had been diagnosed with right-sided clear-cell RCC at age 24 years. His father (now deceased) and paternal uncle had also developed clear cell RCC, at ages 24 and 73 years, respectively. The presence of the c.136C>T mutation in the affected uncle was confirmed subsequently. Case subject 2: A germline heterozygous missense mutation, p.Arg46Gln (c.137G>A, Figure 1) was identified in a 44-year-old woman who presented with bilateral RCC at age 30 years. Case subject 3: A novel germline heterozygous missense mutation,

Affiliations of authors: Cancer Research UK Renal Molecular Oncology Group, Department of Medical and Molecular Genetics, University of Birmingham School of Medicine, Edgbaston, Birmingham, UK (CR, ERW, PK, MRM, DA, FL, ERM); West Midlands Regional Genetics Service, Birmingham Women's Hospital, Birmingham, UK (ERM).

Correspondence to: Eamonn R. Maher, Department of Medical and Molecular Genetics, University of Birmingham, Institute of Biomedical Research, Edgbaston, Birmingham, B15 2TT, UK (e-mail: e.r.maher@bham.ac.uk).

See "Notes" following "References."

DOI: 10.1093/jnci/djn254

© The Author 2008. Published by Oxford University Press. All rights reserved. For Permissions, please e-mail: journals.permissions@oxfordjournals.org.

CONTEXT AND CAVEATS

Prior knowledge

Germline mutations in *SDHB*, a gene encoding a subunit of the mitochondrial enzyme succinate dehydrogenase (SDH), have been found in patients with renal cell carcinoma (RCC) who had a personal or family history of pheochromocytoma or paraganglioma. It was not known whether mutations would be found in genes encoding subunits of SDH in people with evidence of a familial history of RCC but with no family history of these other cancers.

Study design

People for whom there was evidence of increased susceptibility to RCC (ie, family history or early age of diagnosis), but for whom there was no evidence of family history of pheochromocytoma or paraganglioma, were tested for germline mutations in genes encoding three subunits of SDH.

Contribution

Some people with evidence of inherited susceptibility to RCC but for whom there was no other evidence of another cancer susceptibility syndrome harbored germline mutations in *SDHB*.

Implication

Inherited mutations in *SDHB* may predispose individuals to RCC.

Limitations

The added risk of RCC conferred by mutations in *SDHB* and the benefit of screening for these mutations in patients with evidence of familial RCC remain to be determined.

From the Editors

p.Arg11His (c.32G>A, Figure 1) was identified in a 46-year-old man who presented with bilateral eosinophilic chromophobe RCC at age 38 years. p.Arg11 and p.Arg46 are conserved in *SDHB* orthologues from chimpanzee, horse, dog, chicken, mouse, rat, African clawed frog, puffer fish, and drosophila. There was no family history of RCC for case subject 2 or 3, and no other family members were available for mutation testing. None of the four mutation carriers reported a personal or family history of pheochromocytoma or head and neck paragangliomas, although two of the mutations identified (c.136C>T→p.Arg46Stop and c.137G>A→Arg46Gln) have been described previously in individuals with




Figure 1. *SDHB* mutations in probands with familial or bilateral renal cell carcinoma. The mutations detected in the probands are shown in the diagram above with electropherograms of the wild-type sequence above and the mutation below. The nucleotide and amino acid changes are shown (the nucleotide number is based on the A of the ATG start codon being nucleotide 1).

pheochromocytoma and/or head and neck paragangliomas (19–21). None of the three mutations were present in the National Center for Biotechnology Information Single-Nucleotide Polymorphism database (<http://www.ncbi.nlm.nih.gov/projects/SNP/>) or detected in 600 anonymized laboratory control chromosomes from UK population.

Our results show, to our knowledge for the first time, that germline *SDHB* mutations may be present in patients with familial or multicentric RCC without any personal or family history of pheochromocytoma or paraganglioma. Previously, three patients with early-onset RCC were reported in two families with germline *SDHB* mutations (16). Two of these patients had a personal and family history of paraganglioma, and the mother of the third patient was diagnosed with malignant paraganglioma. Subsequently, there were reports of two patients with RCC and a personal or family history of pheochromocytoma/paraganglioma who had germline *SDHB* mutations (22,23). However, *SDHB* mutations have not been considered to be a cause of familial RCC and the risk of RCC was not considered to be elevated in those with *SDHB* mutations.

There are limitations to our study. The study group analyzed was not based on referrals to a specialized center and may not exactly reflect population-based cases of familial RCC. In addition, gene sequencing will not detect all mutations, such as exon deletions or mutations deep within an intron.

Our findings suggest that individuals presenting with features of inherited RCC susceptibility should be screened for germ-

line *SDHB* mutations because surveillance for *SDHB*-related tumors can then be offered to mutation-positive patients and relatives. Many centers undertake routine annual abdominal MRI screening for adrenal and extra-adrenal pheochromocytoma in *SDHB* mutation carriers, and screening for RCC should be considered for inclusion in such surveillance programs. Although histopathologic analysis can inform genetic testing priorities in familial RCC (eg, germline *VHL* mutations are invariably associated with clear-cell RCC and *MET* mutations with papillary RCC), there does not appear to be such a close correlation between germline *SDHB* mutations and histopathologic subtypes of RCC. Although most *SDHB*-associated RCCs are clear-cell RCC, other types of disease may occur (23). All three RCC-associated *SDHB* mutations that we detected were located in the first 50 amino acids. However, it was not possible to deduce clear genotype–phenotype correlations because two of the mutations that we detected have previously been detected in families with pheochromocytoma/paraganglioma without RCC, and RCC has also been associated with *SDHB* mutations closer to the 3' end of the gene (eg, c.847-50delTCTC) (16). More extensive analysis of cohorts of patients with inherited RCC and screening for RCC in *SDHB* patient cohorts will provide information on genotype–phenotype correlations that might influence clinical management.

Pheochromocytomas in patients who carry *VHL* or *SDHB* mutations show evidence of increased expression of HIF-1 and HIF-2 (24). In addition, *SDHB* mutations and missense *VHL* mutations have been

suggested to predispose to pheochromocytoma development by inhibiting normal developmental apoptosis of sympathoadrenal cells via a HIF-independent pathway (25). Genotype-phenotype correlations in VHL disease have closely linked loss of pVHL's ability to inhibit HIF to risk of RCC (26,27), and HIF-2 overexpression can overcome pVHL tumor suppressor activity (28). In contrast to what is observed for *VHL* (5), somatic mutations of *SDHB* appear to be rare in sporadic RCC (18). Further study of the relationship between *SDHB* mutation spectrum, RCC risk, and the functional effect of specific *SDHB* mutations on HIF-1 and HIF-2 expression could provide further insight into the pathogenesis of RCC in individuals with *SDHB* mutations and the role of HIF-dependent and HIF-independent pathways in familial pheochromocytoma and RCC.

References

1. Thoenes W, Storkel S, Rumpelt HJ. Histopathology and classification of renal cell tumors (adenomas, oncocytomas and carcinomas). The basic cytological and histopathological elements and their use for diagnostics. *Path Res Pract*. 1986;181(2):125–143.
2. Maher ER. Von Hippel-Lindau disease. *Curr Mol Med*. 2004;4(8):833–842.
3. Pavlovich CP, Schmidt LS. Searching for the hereditary causes of renal-cell carcinoma. *Nat Rev Cancer*. 2004;4(5):381–393.
4. Linehan WM, Pinto PA, Srinivasan R, et al. Identification of the genes for kidney cancer: opportunity for disease-specific targeted therapeutics. *Clin Cancer Res*. 2007;13:671s–679s.
5. Latif F, Tory K, Gnarr J, et al. Identification of the von Hippel-Lindau disease tumour suppressor gene. *Science*. 1993;260(5112):1317–1320.
6. Schmidt L, Duh F-M, Chen F, et al. Germline and somatic mutations in the tyrosine kinase domain of the MET proto-oncogene in papillary renal carcinomas. *Nat Genet*. 1997;16(1):68–73.
7. Nickerson ML, Warren MB, Toro JR, et al. Mutations in a novel gene lead to kidney tumors, lung wall defects, and benign tumors of the hair follicle in patients with the Birt-Hogg-Dubé syndrome. *Cancer Cell*. 2002;2(2):157–164.
8. Teh B, Giraud S, Sari F, et al. Familial non-VHL non-papillary clear-cell renal cancer. *Lancet*. 1997;349(9055):848–849.
9. Woodward ER, Clifford SC, Astuti D, et al. Familial clear cell renal cell carcinoma (FCRC): clinical features and mutation analysis of the VHL, MET, and CUL2 candidate genes. *J Med Genet*. 2000;37(5):348–353.
10. Tomlinson IP, Alam NA, Rowan AJ, et al. Germline mutations in FH predispose to dominantly inherited uterine fibroids, skin leiomyomata and papillary renal cell cancer. *Nat Genet*. 2002;30(4):406–410.
11. Lehtonen HJ, Kiuru M, Ylisaukko-Oja SK, et al. Increased risk of cancer in patients with fumarate hydratase germline mutation. *J Med Genet*. 2006;43(6):523–526.
12. Baysal BE, Ferrell RE, Willett-Brozick JE, et al. Mutations in SDHD, a mitochondrial complex II gene, in hereditary paraganglioma. *Science*. 2000;287(5454):848–851.
13. Niemann S, Muller U. Mutations in SDHC cause autosomal dominant paraganglioma, type 3. *Nat Genet*. 2000;26(3):268–270.
14. Astuti D, Douglas F, Lennard TW, et al. Germline SDHD mutation in familial pheochromocytoma. *Lancet*. 2001;357:1181–1182.
15. Astuti D, Latif F, Dallol A, et al. Gene mutations in the succinate dehydrogenase subunit SDHB cause susceptibility to familial pheochromocytoma and to familial paraganglioma. *Am J Hum Genet*. 2001;69(1):49–54.
16. Vanharanta S, Buchta M, McWhinney SR, et al. Early-onset renal cell carcinoma as a novel extraparaganglial component of SDHB-associated heritable paraganglioma. *Am J Hum Genet*. 2004;74(1):153–159.
17. Pasini B, McWhinney SR, Bei T, et al. Clinical and molecular genetics of patients with the Carney-Stratakis syndrome and germline mutations of the genes coding for the succinate dehydrogenase subunits SDHB, SDHC, and SDHD. *Eur J Hum Genet*. 2008;16(1):79–88.
18. Morris MR, Maina E, Morgan NV, et al. Molecular genetic analysis of FH-1, FH, and SDHB candidate tumour suppressor genes in renal cell carcinoma. *J Clin Pathol*. 2004;57(7):706–711.
19. Benn DE, Gimenez AP, Reilly JR, et al. Clinical presentation and penetrance of pheochromocytoma/paraganglioma syndromes. *J Clin Endocrinol Metab*. 2006;91(3):827–836.
20. Bayley JP, van Minderhout I, Weiss MM, et al. Mutation analysis of SDHB and SDHC: novel germline mutations in sporadic head and neck paraganglioma and familial paraganglioma and/or pheochromocytoma. *BMC Med Genet*. 2006;7(1):1–10.
21. Gimenez-Roqueplo AP, Favier J, Rustin P, et al. Functional consequences of a SDHB gene mutation in an apparently sporadic pheochromocytoma. *J Clin Endocrinol Metab*. 2002;87(10):4771–4774.
22. Neumann HP, Pawlu C, Peczkowska M, et al. Distinct clinical features of paraganglioma syndromes associated with SDHB and SDHD gene mutations. *JAMA*. 2004;292(8):943–951.
23. Srirangalingam U, Walker L, Khoo B, et al. Clinical manifestations of familial paraganglioma and pheochromocytomas in succinate dehydrogenase B gene mutation carriers [published online ahead of print April 14, 2008]. *Clin Endocrinol*. doi:10.1111/j.1365-2265.2008.03274.x.
24. Pollard PJ, El-Bahrawy M, Poulson R, et al. Expression of HIF-1alpha, HIF-2alpha (EPAS1), and their target genes in paraganglioma and pheochromocytoma with VHL and SDH mutations. *J Clin Endocrinol Metab*. 2006;91(11):4593–4598.
25. Lee S, Nakamura E, Yang H, et al. Neuronal apoptosis linked to EglN3 prolyl hydroxylase and familial pheochromocytoma genes: developmental culling and cancer. *Cancer Cell*. 2005;8(2):155–167.
26. Clifford SC, Cockman ME, Smallwood AC, et al. Contrasting effects on HIF-1alpha regulation by disease-causing pVHL mutations correlate with patterns of tumorigenesis in von Hippel-Lindau disease. *Hum Mol Genet*. 2001;10(10):1029–1038.
27. Li L, Zhang L, Zhang X, et al. Hypoxia-inducible factor linked to differential kidney cancer risk seen with type 2A and type 2B VHL mutations. *Mol Cell Biol*. 2007;27(15):5381–5392.
28. Kondo K, Kim WY, Lechpammer M, et al. Inhibition of HIF2alpha is sufficient to suppress pVHL-defective tumor growth. *PLoS Biol*. 2003;1(3):E83.

Notes

We thank Cancer Research, United Kingdom, for financial support and the many clinician and patients who helped with this work. The authors take full responsibility for the study design, data collection, analysis and interpretation of the data, the decision to submit the manuscript for publication, and the writing of the manuscript.

Manuscript received February 8, 2008; revised June 10, 2008; accepted June 19, 2008.