

The Immunotherapy and Gene Therapy of Cancer

By Steven A. Rosenberg

IN THE LAST decade biologic therapy has emerged as a mode of treatment capable of mediating the regression of cancer in some patients. Biologic therapy differs conceptually from surgery, radiation therapy, or chemotherapy because it acts, not by directly attacking the tumor, but by stimulating natural host defense mechanisms to mediate cancer regression.^{1,2} The predominant natural host defense mechanism is the immune system.

Two factors have converged to enable new approaches to the immune manipulation of the cancer patient in an effort to mediate tumor regression. Substantial progress has been made in understanding the molecular nature of the cellular immune response to cell-surface antigens or to exogenous antigens presented in conjunction with major histocompatibility complex (MHC) molecules. In addition, rapid advances in biotechnology have made it possible to obtain large amounts of biologic molecules that were previously available only in minute quantities. Most prominent among these scarce biologic molecules are the cytokines, which are hormones produced by immune cells that play important roles in regulating the immune system.


We have attempted to develop methods for mediating the rejection of human cancers using immune manipulations with recombinant cytokines and immune cells. One approach we have taken has been referred to as adoptive immunotherapy, which can be defined as the transfer, to the tumor-bearing host, of immunologically reactive cells with antitumor reactivity that can mediate antitumor effects either directly or indirectly. The inability to identify immune cells in the tumor-bearing animal and human that could specifically recognize tumor cells represented the major impediment to the development of adoptive immunotherapy.³

The description in 1976 of a hormone called T-cell growth factor (later renamed interleukin-2 [IL-2]) provided new opportunities for identifying cells with antitumor reactivity from a cancer-bearing patient.⁴ IL-2 plays

a central role in regulating the immune response by mediating the expansion of lymphocyte clones responsive to antigen *in vivo*.⁵

STUDIES OF IL-2 AND LYMPHOKINE-ACTIVATED KILLER CELLS

Studies with IL-2 attempting to isolate lymphocytes with the specific ability to recognize cancer but not normal cells led us to the description of lymphokine-activated killer (LAK) cells in 1980.^{6,7} The incubation of lymphocytes in IL-2 results in the generation of cells capable of lysing fresh tumor cells in short-term, ie, 4-hour, cytotoxicity assays. We performed extensive studies of the biologic characteristics of these LAK cells in both the mouse and human and identified them as non-B, non-T cells that were capable of recognizing cancer cells in a non-MHC-restricted fashion.⁸⁻¹⁸ LAK cells, which could be generated from either the normal or tumor-bearing host, appeared to represent a primitive immunosurveillance system capable of recognizing and destroying altered cells. Extensive studies in experimental animal models subsequently demonstrated that the administration of either IL-2 alone or LAK cells plus IL-2 could mediate the regression of established lung and liver metastases from a variety of animal cancers.¹⁹⁻²⁹ The antitumor effects of both IL-2 and LAK cells were dose-related, and in most models, the combined admin-



This lecture was delivered to the American Society of Clinical Oncology on May 20, 1991.

Address reprint requests to Steven A. Rosenberg, MD, Chief of Surgery, National Cancer Institute, National Institutes of Health, Bethesda, MD 20892.

© 1992 by American Society of Clinical Oncology.
0732-183X/92/1002-0021\$3.00/0




Fig 1. Therapy of established liver metastases in mice using either recombinant IL-2 alone (●) or LAK cells plus recombinant IL-2 (○). Varying doses of recombinant IL-2 were administered intraperitoneally every 8 hours for 5 days. At each dose of recombinant IL-2 more effective reduction in established 3-day liver metastases was seen when LAK cells were also administered. Reprinted with permission from Lafreniere et al.²⁴

istration of LAK cells plus IL-2 mediated better antitumor effects than IL-2 alone. An example of one of these early studies in the treatment of established murine liver metastases is shown in Fig 1.²⁴ A summary of the major findings of the use of either IL-2 alone or LAK cells plus

IL-2 in animal therapy models is shown in Tables 1 and 2. These early studies demonstrated that LAK cells could expand in vivo under the influence of their requisite growth factor, IL-2, and maintain their antitumor activity as they multiplied. This basic principle of adoptive immunotherapy influenced many of our subsequent treatments. Activated cells could expand in vivo as long as IL-2 was administered and died in vivo when IL-2 administration was discontinued.^{30,31}

The application of these findings to patients with advanced cancer was dependent on a supply of large amounts of IL-2, a problem that was solved when the gene for IL-2 was cloned and expressed in *Escherichia coli*.^{32,33} Purification to homogeneity of this recombinant IL-2 and the demonstration that it was similar in all biologic and functional aspects to natural IL-2^{34,35} led directly to clinical trials in patients with advanced cancer in whom all available effective treatments had failed.

Although results of studies in animal models predicted that the combined administration of LAK cells and IL-2 would mediate greater antitumor effects than either treatment alone, neither IL-2 nor activated killer cells had been administered to cancer patients. We began phase I studies of the administration of natural Jurkat-cell-derived IL-2 and, later, recombinant IL-2 to explore different doses, routes, and schedules of administration.^{36,37} These studies were conducted in 39 patients with advanced cancer, none of whom responded to

Table 1. Immunotherapy of Murine Tumors With IL-2 Alone

1. Liver and lung micrometastases (3-day) from a variety of immunogenic and nonimmunogenic sarcomas, melanomas, and adenocarcinomas can be inhibited by IL-2 administration.
2. Lung macrometastases (10-day) from two immunogenic sarcomas (but not from two nonimmunogenic sarcomas) can be inhibited by IL-2 administration.
3. A direct relationship exists between the dosage of IL-2 and therapeutic effect.
4. High-dose IL-2 administration leads to in vivo lymphoid proliferation in visceral organs. These cells have LAK activity in vitro.
5. The immunotherapeutic effect of IL-2 on 3-day micrometastases is mediated by asialo GM1⁺ LAK cells. In immunogenic tumors, Lyt 2⁺ cells also participate.
6. The immunotherapeutic effect of IL-2 on 10-day macrometastases is mediated by Lyt 2⁺ cells.
7. Immunosuppression with irradiation or cyclophosphamide can inhibit IL-2 therapy against 3-day metastases, but can enhance the effects of IL-2 on 10-day macrometastases.
8. The sensitivity of macrometastases to therapy involving IL-2 appears to be directly related to the expression of MHC antigens (class I) on the tumor.
9. The administration of IL-2 can enhance the therapeutic effect of concomitantly administered LAK cells, TIL, and, specifically, sensitized T lymphocytes.

Abbreviation: TIL, tumor-infiltrating lymphocytes.

Table 2. Immunotherapy of Murine Tumors With LAK Cells Plus IL-2

1. Liver and lung micrometastases (3-day) from a variety of immunogenic and nonimmunogenic sarcomas, melanomas, and adenocarcinomas can be inhibited by treatment with LAK cells plus IL-2.
2. A direct relationship exists between therapeutic effect and the dosage of IL-2 and the dose of LAK cells.
3. The precursor of the LAK cell effective in vivo is Thy 1-Ig-a-ASGM1.
4. Three-day incubation of splenocytes is optimal for the generation of LAK cells effective in vivo.
5. Immunotherapy of micrometastases with LAK cells and IL-2 is effective in hosts suppressed by total body irradiation or treatment with cyclophosphamide. Therapy is also effective in B mice (thymectomized, lethally irradiated, reconstituted with T-cell-depleted bone marrow).
6. Immunotherapy of micrometastases with allogeneic LAK cells plus IL-2 is effective.
7. LAK cells effective in immunotherapy can be generated from the splenocytes of tumor-bearing mice.
8. Metastases that persist after in vivo therapy with LAK cells plus IL-2 are sensitive to LAK-cell lysis both in vitro and in subsequent in vivo experiments. We have been unable to generate LAK-resistant tumor cells.
9. Administration of IL-2 leads to in vivo proliferation of transferred LAK cells.
10. Diffuse intraperitoneal carcinomatosis can be successfully treated with intraperitoneal LAK cells plus IL-2.
11. LAK cells can mediate antibody-dependent cellular cytotoxicity, and this administration of IL-2 alone or LAK cells plus IL-2 can enhance the in vivo therapeutic efficacy of monoclonal antibodies with antitumor reactivity.

treatment.³⁸ Similarly, studies were conducted with activated killer cells, first using phytohemagglutinin-activated killer (PAK) cells and then cells activated with recombinant IL-2 (LAK cells) in 27 cancer patients, and none responded to treatment.

After presenting this information to the Food and Drug Administration, we received permission to treat

patients with a combination of LAK cells and IL-2. These studies were initiated in November 1984. We had previously demonstrated that the administration of IL-2 led to a rapid decrease in LAK cell precursors in the circulation, which rebounded substantially after discontinuing IL-2 administration.^{36,37} We developed a treatment protocol in which IL-2 was administered for one cycle followed by repeated leukophereses and subsequent administration of LAK cells along with IL-2. IL-2 was administered as an intravenous bolus injection every 8 hours. Antitumor effects in cancer patients were seen with this combined treatment.³⁹ As we gained confidence in the administration of IL-2 and the management of its side effects, we used higher doses of IL-2 alone and began to see antitumor effects with this treatment as well.⁴⁰⁻⁴²

The results of the treatment of 320 consecutive patients with advanced cancer treated with either IL-2 alone (140 patients) or IL-2 plus LAK cells (180 patients) are listed in Table 3. These data demonstrate that treatment with LAK cells plus IL-2 can mediate objective regressions in patients with kidney cancer, melanoma, colorectal cancer, and non-Hodgkin's lymphoma. Approximately 10% of patients with renal cell cancer and melanoma achieve complete regression of their advanced metastatic disease. Responses to IL-2 alone have also been seen in patients with kidney cancer and melanoma. Few patients with other histologies have been treated, and thus, definitive comments about response rates in other diseases cannot yet be made. The durations of responses in these patients as of March 1991 are shown in Table 4. Of 20 patients who achieved a complete response, nine remain continuously free of disease from 13 to 75 months.

The ability to achieve antitumor responses with both

Table 3. Immunotherapy of Patients With Advanced Cancer Using IL-2 Alone or With LAK Cells

Diagnosis	IL-2				IL-2 + LAK Cells			
	Assessable*	CR	PR	CR + PR (%)	Assessable†	CR	PR	CR + PR (%)
Renal cell cancer	60	5	6	18	72	8	17	35
Melanoma	41	1	9	24	48	4	6	21
Colorectal cancer	12	0	0	0	30	1	3	13
Non-Hodgkin's lymphoma	11	0	0	0	7	1	3	57
Breast cancer	3	0	0	0	1	0	0	0
Sarcoma	1	0	0	0	6	0	0	0
Lung cancer	1	0	0	0	5	0	0	0
Other	7‡	0	0	0	9§	0	0	0
Total	136	6	15	15	178	14	29	24

*Includes all treated patients except four who died of therapy.

†Includes all treated patients except one lost to follow-up and one dead due to therapy.

‡Two patients each with hepatoma and brain cancer; one each with ovarian, pancreatic, and uterine cancer.

§One patient each with cancer of brain, esophagus, ovary, testes, thyroid, gastrinoma, unknown primary; and two patients with Hodgkin's lymphoma.

Table 4. Duration of Responses (as of 3/91)

Diagnosis	LAK + IL-2		IL-2	
	CR (months)	PR (months)	CR (months)	PR (months)
Renal cell cancer	51+, 47+, 27, 15, 13, 11, 9, 6	45+, 38, 19, 13, 11, 11, 9, 7, 7, 6, 6, 6, 6, 3, 2, 1, 1	51+, 49+, 39+, 38, 36	49+, 43+, 39, 34, 26, 10
Melanoma	75+, 56+, 42+, 13	41, 6, 6, 3, 2, 2	13+	64+, 15, 10, 8, 7, 7, 5, 3, 2
Colorectal cancer	21	11, 6, 6, 2	—	—
Non-Hodgkin's lymphoma	10	55+, 31+, 14	—	—

NOTE. Of 20 patients with CRs, nine remain in CR at 13 to 75 months.

Abbreviations: CR, complete response; PR, partial response.




Fig 2. Results of a prospective randomized trial in patients with advanced melanoma who received either IL-2 alone (○) or IL-2 plus LAK cells (□). Patients receiving IL-2 plus LAK cells had a trend toward improved survival ($P_2 = .062$).




Fig 3. Results of a prospective randomized trial in patients with advanced renal cell cancer who received either IL-2 alone (○) or LAK cells plus IL-2 (□). No difference in long-term survival was seen ($P_2 = .55$).

IL-2 alone and LAK cells plus IL-2 led us to perform a prospective randomized trial in 181 patients with advanced cancer, the majority of whom had renal cell cancer and melanoma. The survival curves with a median follow-up of 50.2 months are shown in Figs 2 and 3. An example of some of the responses seen in patients treated with LAK cells and IL-2 or IL-2 alone are shown in Figs 4 through 6. Treatment with LAK cells and IL-2 appeared to yield improved survival compared with IL-2 alone in patients with advanced melanoma ($P_2 = .06$).




Fig 4. Complete regression of bony metastasis from renal cell cancer in the pubic ramus of a patient treated with LAK cells plus IL-2. This patient underwent complete regression of bony and lung metastases and remains disease-free over 4 years after treatment.




Fig 5. Regression of lung metastases in a patient with advanced malignant melanoma treated with recombinant IL-2 alone.




Fig 6. Objective regression of liver and splenic metastases in the same patient shown in Fig 5.

but no difference in patients with kidney cancer ($P_2 = .55$).

The mechanism of action of these antitumor responses in humans appears to be similar to that seen in the mouse. After administration of this immunotherapy, extensive lymphoid infiltrates could be seen in tumors in conjunction with tumor destruction. An example of a subcutaneous melanoma lesion biopsied before and after treatment with LAK cells and IL-2 is shown in Fig 7.

The administration of large doses of recombinant IL-2 can be associated with substantial side effects. The major toxicity of IL-2 appears to result from an increased membrane permeability induced by IL-2 that leads to fluid and colloid loss into visceral organs and soft tissues.⁴³ These side effects and their mechanisms have been extensively studied and reported.^{42,44-62} A summary of the side effects of treatment with 1,039 treatment courses in our first 652 patients treated with a variety of regimens involving high-dose IL-2, either alone, with activated killer cells, or with other cytokines is shown in Table 5. Treatment-related mortality for the




Fig 7. The histologic appearance of subcutaneous melanoma deposits in a patient being treated with LAK cells and IL-2: (A) pretreatment, (B) posttreatment. There is an extensive infiltrate of lymphocytes and evidence of tumor destruction after treatment.

Table 5. Toxicity of Treatment With IL-2

No. of patients	652
No. of courses	1,039
Chills	399
Pruritus	180
Necrosis	5
Anaphylaxis	1
Mucositis (requiring liquid diet)	30
Alimentation not possible	4
Nausea and vomiting	666
Diarrhea	596
Hyperbilirubinemia, maximum (mg/dL)	
2.1-6.0	547
6.1-10.0	179
10.1+	83
Oliguria	
< 80 mL/8 hours	347
< 240 mL/24 hours	42
Weight gain (% body weight)	
0.0-5.0	377
5.1-10.0	436
10.1-15.0	175
15.1-20.0	38
20.1+	13
Elevated creatinine, maximum (mg/dL)	
2.1-6.0	637
6.1-10.0	85
10.1+	10
Hematuria (gross)	2
Edema (symptomatic nerve or vessel compression)	17
Tissue ischemia	2
Respiratory distress	
Not intubated	67
Intubated	41
Bronchospasm	9
Pleural effusion (requiring thoracentesis)	17
Somnolence	114
Coma	33
Disorientation	215
Hypotension (requiring pressors)	508
Angina	22
Myocardial infarction	6
Arrhythmias	78
Anemia requiring transfusion (no. of units transfused)	
1-5	377
6-10	95
11-15	24
16+	14
Thrombocytopenia, minimum (/μL)	
< 20,000	131
20,001-60,000	361
60,001-100,000	285
Central line sepsis	63
Death	10

entire series was 1.5% of patients. These data do not reflect the first 66 patients treated in the phase I studies, in which no fatalities occurred. As we have gained experience with the administration of high-dose IL-2,

treatment-related morbidity and mortality has decreased, and we have not had a treatment-related fatality in the last 383 patients treated with high-dose IL-2. Shortly after our results were published, the National Cancer Institute (NCI) initiated studies by the IL-2 Working Group using LAK cells and IL-2. Similar treatment results were obtained, and one treatment-related mortality was experienced in the first 94 patients treated.⁶³ A more recent report of 515 patients treated by this working group with a variety of high-dose IL-2-related therapies showed a 3.5% treatment-related mortality. High-dose IL-2 can be administered safely either alone or with activated killer cells, although experience is required with the unique nature of the toxicities resulting from the administration of this biologic agent. A variety of other groups have confirmed the antitumor activity of IL-2 alone and LAK cells plus IL-2.⁶³⁻⁷⁶

STUDIES OF TUMOR-INFILTRATING LYMPHOCYTES

Our studies with LAK cells and IL-2 led us to search for immune cells with improved therapeutic effectiveness. In 1986, we described a class of T lymphocytes with unique antitumor activity that we called tumor-infiltrating lymphocytes (TIL).^{77,78} TIL are cells that infiltrate into growing tumors and can be grown by culturing single-cell suspensions obtained from tumors in IL-2. Although lymphocytes comprise only a small subpopulation of the cells in a cancer nodule, some of these lymphocytes contain IL-2 receptors—presumably because of their interactions with tumor antigens—and grow under the influence of IL-2. Although tumor cells also grow in the culture, the lymphocytes capable of eliminating the tumor cells have a selective growth advantage. After 2 to 3 weeks of culture, pure populations of lymphocytes without contaminating tumor cells are obtained.

In vitro, TIL obtained from methylcholanthrene-induced murine sarcomas, as well as from the MC-38 colon adenocarcinoma, specifically lyse the tumor from which they are derived and not other tumors.⁷⁹ More recently, we have demonstrated specific cytokine release when these TIL are in contact with their autologous tumors.^{80,81} TIL are thus unique in their ability to identify cancer-associated antigens in vitro.

Extensive in vivo studies demonstrated that TIL were from 50 to 100 times more effective in treating established 3-day lung and liver metastases than were LAK cells.⁷⁷ An example of the titration of the antitumor effectiveness of TIL and LAK cells is shown in Fig 8. The results of extensive studies of the mechanism of this




Fig 8. Titration of LAK cells and TIL administered with IL-2 in mice bearing 3-day lung metastases from a transplantable sarcoma⁷⁷⁻⁷⁸: ●, TIL, one treatment; ○, LAK cells, one treatment; ■, LAK cells, two treatments. TIL appear to be 50 to 100 times more potent than LAK cells in reducing established lung metastases. Reprinted with permission from Rosenberg et al⁷⁷ (*Science* 233:1318-1321, 1986). Copyright 1986 by the AAAS.

immunotherapeutic effect of TIL and IL-2 in mice with established tumors are listed in Table 6.⁸⁰⁻⁸⁹ TIL are classic T cells. In the mouse they are almost exclusively CD8+, but human TIL can be both CD4+ and CD8+. We have recently shown that the specific secretion of interferon gamma by TIL cocultured with tumor appears to be the best in vitro correlate of the in vivo antitumor effectiveness of TIL.⁸⁰ Non-lytic TIL that can specifically secrete interferon gamma when in contact with the autologous tumor are effective in treating established lung micrometastases. In recent studies, we have shown that the administration of antibodies to interferon gamma or tumor necrosis factor (TNF) can abrogate some of the therapeutic effectiveness of TIL, thus suggesting a very potent role for cytokine secretion in this phenomenon.⁸⁰

We and others have successfully grown TIL from approximately 80% of over 300 human cancers, including melanoma, renal cell cancer, colon cancer, breast cancer, bladder cancer, neuroblastoma, lymphomas, and other tumor types.⁹⁰⁻¹⁰⁷ From approximately one third of patients with melanoma, TIL can be derived that have the ability to lyse specifically the autologous melanoma and not other melanomas or normal cells from the same patient.⁹⁰⁻⁹² An example of this phenomena is shown in Fig 9.⁹⁰ More recently, we have demonstrated that TIL can also recognize unique tumor antigens on the tumors of some patients with melanoma, breast cancer, colon

cancer, and lymphomas as evidenced by the secretion of specific cytokines when cocultured with the autologous tumor.¹⁰⁸ An example of the specific recognition of breast cancer antigens by this technique is shown in Fig 10. Extensive studies of the in vitro characteristics of human TIL have been performed (Table 7).

The ability of TIL to recognize unique cancer-associated antigens has made the study of the biologic characteristics of these determinants possible. A study of the lytic recognition of a variety of HLA-type melanomas by TIL demonstrated that melanomas from patients that share MHC antigens are often cross-recognized by TIL, as would be expected for an MHC-restricted T-cell reaction.^{109,110} The example of this phenomenon shown in Table 8 demonstrates the recognition of a melanoma antigen restricted by the HLA-A24 class I determinant. TIL from melanoma patient no. 888 recognized autologous tumor and all three allogeneic melanomas from patients bearing the HLA-24 determinant, but did not recognize a normal Epstein-Barr virus (EBV)-transformed line from an HLA-24-positive patient or 16 other tumors that were unmatched at any HLA locus.

In these studies we found that the HLA-A2 class I determinant was frequently used as an MHC restriction element in the recognition of melanoma antigens. To study this phenomenon further, the HLA-A2 gene was transfected into the allogeneic melanomas of six dif-

Table 6. Immunotherapy of Murine Tumors With TIL and IL-2

1. The administration of TIL plus IL-2 is 50 to 100 times more effective than LAK cells plus IL-2 in reducing established (day 3) lung micrometastases.
2. The administration of TIL plus IL-2 plus cyclophosphamide is effective in treating mice with advanced lung (day 14) or liver (day 8) metastases. All three agents are required for effective treatment.
3. The phenotype of TIL effective in vivo is CD3+, CD4-, CD8+.
4. TIL raised from tumors that express class I antigens only after transfection with genes coding for class I can mediate the regression of micrometastases from the parental tumor after treatment of the mouse with interferon gamma.
5. Local tumor irradiation synergizes with TIL and IL-2 administration in mediating the regression of established metastases. Local irradiation can substitute for cyclophosphamide.
6. TIL with improved antitumor activity in vivo can be generated from tumor suspensions by using immunomagnetic beads to isolate lymphocytes followed by incubation of lymphocytes in low-dose IL-2.
7. In mice cured of lung micrometastases by administration of TIL, live TIL can be identified in vivo 3 months after injection.
8. The specific secretion of interferon gamma by TIL when cultured with tumor is the best in vitro correlate of the in vivo antitumor effectiveness of TIL. Nonlytic TIL that specifically secrete interferon gamma can effectively treat established lung micrometastases.




Fig 9. Specific TIL lysis of autologous melanoma and not other allogeneic melanomas. TIL from three patients (○, patient A; □, patient B; ●, patient C) were tested simultaneously against melanoma from these patients. Only the autologous tumors exhibited significant lysis. Reprinted with permission from Muul et al.⁹⁰

ferent HLA-A2-negative patients, and HLA-A2-expressing and -nonexpressing clones of these melanomas were isolated and tested for lysis by an HLA-A2-restricted melanoma TIL. The allogeneic melanomas of all six patients shared tumor antigens uncovered by insertion of the HLA-A2 normal class I gene. Thus, it appears that human melanomas bear at least some antigens that are commonly expressed between individuals and are recognized in an MHC-restricted fashion.

The presence of shared tumor antigens provides the possibility that these antigens might be used for active

immunization of patients against their own tumors. Libraries of cDNA have been prepared from human melanomas and are being transfected into nonantigen-expressing autologous lines or transformed fibroblast lines in an attempt to identify the antigen recognized by TIL. The identification of such cancer antigen genes might enable the insertion of the gene into viruses for use in the immunization against cancer in patients with advanced disease or for the primary or secondary prevention of cancer. These efforts are being actively pursued in the laboratory.

The increased efficacy of murine TIL in treating established cancers in mice and the ability to generate TIL from a variety of human tumors led to a pilot trial of




Fig 10. Demonstration of unique tumor antigens on breast cancer as evidenced by specific cytokine secretion by TIL. TIL from a patient with breast cancer secreted granulocyte-macrophage colony-stimulating factor, TNF alpha, and interferon gamma only when cocultured with the autologous breast cancer (●) and not with an autologous B-cell line (△) or with four other allogeneic breast cancers: □, ▽, ◇, and ⊙. Reprinted with permission from Schwartzentruber et al.¹⁰⁸

Table 7. In Vitro Studies of Human TIL

1. TIL can be grown in IL-2 from approximately 80% of human cancers of a variety of histologic types including melanoma, renal cell cancer, colon cancer, breast cancer, ovarian cancer, and others.
2. TIL with specific cytolytic activity for fresh cancer cells can be grown from approximately one third of patients with melanoma.
3. TIL are mainly CD3+ and can be either CD8+, CD4+, or mixtures of both. Smaller numbers of CD56+ TIL can also be present.
4. Specific lysis of TIL can be inhibited by antibodies to CD3 or to MHC class I molecules.
5. Growth of TIL from melanoma patients in IL-2 plus IL-4 results in increased in vitro lytic specificity for autologous melanoma.
6. Incubation of cultured tumors in interferon gamma increases their susceptibility to lysis by TIL.
7. TIL with lytic specificity for autologous tumor have not been obtained from patients with colorectal cancer, breast cancer, or sarcomas and only rarely from patients with renal cell cancer.
8. Direct positive panning techniques using antibody-coated flasks can be used to separate and grow highly purified subpopulations of CD4+ and CD8+ TIL.
9. Repeated immunoselection using TIL with specific lysis can be used to identify tumor lines resistant to lysis by autologous TIL. These immunoselected tumor can be used to identify multiple tumor antigens on a single tumor.
10. Shared tumor antigens on allogeneic melanomas that are recognized in an MHC-restricted fashion can be identified by testing lysis of specific TIL on panels of HLA-type melanoma cultures. HLA-A2 is a common restriction element in the recognition of melanoma antigens.
11. MHC-restricted recognition of shared melanoma antigens on allogeneic melanomas was demonstrated by using HLA-A2-restricted specific TIL to lyse allogeneic melanomas transfected with the gene for HLA-A2.
12. Nonlytic TIL with specific immune recognition of human tumor antigens can be recognized by the specific release of cytokines (such as granulocyte-macrophage colony-stimulating factor, TNF, and interferon gamma) after incubation with autologous tumor. Specific reactivity has been identified in patients with melanoma and breast cancer.
13. In vitro lysis of autologous tumor by TIL exhibits mild but significant correlation with clinical response.

TIL therapy in 12 patients with advanced cancer.¹¹¹ In prior murine adoptive immunotherapy models in which bulk tumor was treated, immunosuppression with either cyclophosphamide or total body irradiation was necessary to achieve optimal therapeutic effects.^{112,113} Thus, in the pilot trial, patients received TIL with varying doses of IL-2 with or without cyclophosphamide. After this pilot trial, a phase II trial was initiated in patients with metastatic melanoma using TIL with high-dose IL-2.¹¹⁴ The results of this trial are shown in Table 9. Response rates of 36% to 39% were seen in patients in whom IL-2-containing regimens had previously failed. In preliminary results, it does not appear that the administra-

tion of cyclophosphamide is required to achieve the therapeutic effects of TIL and IL-2, although a randomized trial will be needed to confirm this.

In conjunction with these trials, studies were conducted to determine whether TIL accumulate in tumor deposits. TIL were labeled with indium-111 and infused along with the administration of IL-2.^{115,116} Clear tumor localization was seen in 13 of 18 radionuclide scan series. In addition, biopsies of tumors and normal skin showed a slow accumulation of indium-111 TIL in cancer deposits compared with normal tissue (Fig 11). These studies showed that TIL recirculate and specifically localize in cancer deposits and provided a basis for future experiments using gene-modified TIL in the treatment of human cancer.

After these early studies with IL-2, LAK cells, and TIL, attempts were made to find predictors of antitumor response. An analysis of responding and nonresponding lesions to IL-2-based immunotherapies in humans suggested a direct relationship between the expression of class II antigens on the tumor-cell surface and the response of individual lesions.¹¹⁷ These observations led to experimental studies in mice using IL-2 combined with interferon alfa to upregulate MHC antigens on tumor cells.¹¹⁸ Thus, tumors that were relatively resistant to treatment with IL-2 or interferon alone were treated with the combination of these two agents at a variety of doses. Only the combination of IL-2 and interferon alfa was capable of reducing the number of established metastases in these mice. Further studies showed that the addition of TIL to IL-2 and interferon alfa increased antitumor effectiveness compared with the use of IL-2 and interferon alone or the use of TIL and IL-2 alone.

A phase I study was undertaken in 201 humans with advanced cancer who received escalating doses of IL-2 and interferon alfa that suggested that the combination might be more effective in the treatment of patients with advanced kidney cancer and melanoma than either of these agents alone¹¹⁹ (Table 10). A conclusive test of this hypothesis, however, would require a prospective randomized trial.

A variety of experimental leads for improving immunotherapy with IL-2 or TIL are listed in Table 11. The combination of TIL plus local radiation therapy has shown synergistic activity against localized tumor deposits in mice.⁸³ Possible methods for generating TIL with improved effectiveness involve the use of lymphocyte subpopulations or clones or restimulation with tumor in vitro. TIL with enhanced in vitro lytic specificity have been raised in humans using IL-2 in combination with interleukin-4 (IL-4).¹²⁰ Recently, Yang et al have shown

Table 8. Cytolytic Activity of Patient No. 888 TIL Against HLA-Matched and Unmatched Targets

% Lysis*	Autologous	Unmatched	HLA-A1†	HLA-A24	HLA-B22	HLA-B52	HLA-Cw1	HLA-Cw7
> 10	888; 29 ± 2; (6) 888-mel; 46 ± 3; (14)			878; 12 ± 1; (7) 809; 14 ± 1; (6) 501-mel; 16 ± 1; (9)				(501)
≤ 10	888-EBV; -8 ± 5 888, fibroblast 1	894; 2 ± 0 890; 0 889; 0 865; 2 851‡; -1 ± 1 822; 2 ± 1; (6) 816; 1 796; -1 745; 6 734; 1 697; 0 ± 1 583-EBV; -4 ± 2 553; 2 526-mel; -1 ± 0; (7) Z2B; -1 Daudi; -1 ± 1; (17)	790; 1 537-mel; -2 ± 1; (6) 537-EBV; -4 ± 2; (3) 397-mel; -1 ± 1; (6)	501-EBV; -2 ± 0	544-EBV; -3		(537-mel) (537-EBV)	890; 1 ± 1; (3) 882; 0 ± 1 864; 3 ± 2; (3) 836; 2 ± 1; (4) 619; 0 624-mel; -1 ± 2; (4) 586-mel; -1 ± 1; (11) 586-EBV; -12 ± 7 551-EBV; -2 ± 5 (790) (501-EBV)

NOTE. Data indicate target cells; mean % lysis ± SEM; (n), which is 1 where no SEM is indicated or 2 unless otherwise specified. Target cells are fresh cryopreserved melanomas unless otherwise indicated. Targets matching more than one TIL HLA locus are indicated in parentheses.

Abbreviations: mel, cultured melanoma line; EBV, EBV-transformed B-cell line; ⁵¹Cr, radioactive isotope of chromium; Ag, antigen.

*Lysis in 4-hour ⁵¹Cr-release assays at effector:target ratio of 40:1. Values ≤ 10% are not considered significant.

†Target donor HLA Ag matching TIL effector HLA Ag.

‡Fresh sarcoma target.

that murine TIL with far greater in vivo efficacy can be generated using low-dose IL-2 (ie, 10 to 20 U/mL) compared with the higher-dose IL-2 currently in use (1,000 U/mL).⁸⁴ Each of these modifications in TIL preparation requires evaluation as it may lead to the generation of TIL with improved therapeutic effectiveness in humans.

Cytokine hormones are being described with increasing frequency, and many have potent immune-stimulating activity. We have recently demonstrated that the administration of interleukin-6 (IL-6) can successfully treat lung micrometastases in animal models.¹²¹ Interleu-

kin-7 (IL-7) can be used to generate cells with antitumor activity from draining lymph nodes in mice.¹²² These agents warrant evaluation in humans with advanced cancer. The combined use of monoclonal antibodies and LAK cells, which mediate potent antibody-dependent cellular cytotoxicity, leads to increased therapeutic effect in murine models.¹²³⁻¹²⁵ Finally, studies combining these immunotherapy approaches with chemotherapy regimens¹²⁶ or after surgery to reduce tumor burden represent alternate approaches worthy of investigation in humans.

GENE THERAPY FOR CANCER

Attempts to alter immune reactivity in the cancer-bearing host have recently turned to genetic manipulation of immune cells and tumor cells.¹²⁷ Gene therapy can be defined as a therapeutic technique in which a functioning gene is inserted into the cells of the patient to correct an inborn genetic error or to provide a new function to the cell. The development of techniques for inserting and expressing foreign genes in eukaryotic cells and an increased understanding of the regulation of gene expression have opened new possibilities for cancer therapy based on these gene transfer techniques.

Our previous studies demonstrating that TIL accumu-

Table 9. TIL Treatment of Patients With Melanoma

	NR	Objective Response (PR + CR)	PR + CR (all)	%
No previous IL-2				
IL-2 + CY	17	11	11/28	39
IL-2 (no CY)	7	4	4/11	36
Previous IL-2				
IL-2 + CY*	3	3	3/6	50
IL-2 (no CY)	4	1	1/5	20

NOTE. Excludes patients with brain metastases at start of treatment (all NR).

Abbreviations: NR, no response; CY, cyclophosphamide.

*Excludes two patients who received IL-2 at 30,000/kg; both were NR.




Fig 11. After intravenous injection, TIL accumulate in tumor deposits. Indium-111-labeled TIL were administered to patients (○, ●, △, ▲), and normal tissue (A) and tumors (B) were biopsied. There was no accumulation in normal tissue, but a slow accumulation of labeled TIL was seen in the tumor of these four patients. Reprinted with permission from Fisher et al.¹¹⁵

lated in tumor deposits suggested that TIL could be used as vehicles to deliver to the tumor-site molecules that would increase their antitumor effectiveness.^{115,116} Using gene transfer techniques, new properties could be introduced into TIL to enhance existing functions or to generate a kind of lymphocyte that normally did not exist in nature. Along with my colleagues (W. French Anderson, National Heart, Lung and Blood Institute; R. Michael Blaese, NCI), we have begun to study genetic modification of TIL in an effort to increase their therapeutic efficacy.

These studies were designed in two phases. The first

was an attempt to insert a gene coding for neomycin phosphotransferase that would mark these cells with a bacterial enzyme that would render them resistant to the antibiotic neomycin and, thus, enable us to perform studies of the long-term distribution and survival of TIL in humans. These studies were conceived as a preliminary effort to demonstrate the safety and efficacy of using retroviral-mediated gene transduction to introduce genes into TIL and as a preparation for introducing genes that could improve TIL therapeutic efficacy. Although most physical techniques for introducing genes into the DNA of eukaryotic cells are very inefficient and at best can introduce genes into only one in 10,000 cells, the use of retroviral vectors can introduce genes with an efficiency up to 30% to 50%.¹²⁷⁻¹³⁰ After infection of the cell, viral RNA is transcribed to DNA, which randomly integrates into the genome. The integrated provirus is indistinguishable from other cellular genes and replicates as does normal DNA during mitosis. In the early 1980s techniques were developed for generating replication-incompetent retroviruses using packaging cell lines. These cell lines can produce genetically modified virus lacking the genes for virally encoded proteins so that the retrovirus can infect subsequent cells but cannot itself form new virus. The retroviral vector that we used was derived from a murine Moloney leukemia virus genetically engineered to contain the gene for neomycin phosphotransferase.^{131,132}

A potential danger of using a retrovirus to insert genes into human cells was the possibility that replication-competent virus might be produced by recombination of vector sequences with viral coding sequences present in the packaging cell line. Careful modification of the packaging cell lines and the retroviral vectors were performed to minimize the possibility of this helper virus generation.^{131,132} The random incorporation of the retroviral-coded DNA into the genome could also activate sequences in the transduced cells that could result in autonomous growth, and this required careful evaluation before reinfusion of cells into patients.

The use of this neomycin phosphotransferase marker gene in TIL provided many advantages that did not exist using the indium-111 label. Indium-111 has a half-life of 2.8 days and the combination of the natural decay of the isotope and the spontaneous release of the label from the cell severely limited the time that these cells could be used to study cell survival in vivo. Autoirradiation of the cell resulting from indium-111 incorporation led to potential damage of TIL that could alter their function. Neomycin resistance genes appeared to represent an ideal label that would be incorporated into the genome of the cell without altering its function and would

Table 10. Treatment of Patients With Advanced Cancer Using Interferon alfa Plus IL-2

IFN (U/m ²)	IL-2 (×10 ⁶)	Melanoma				Renal Cell Cancer				Colorectal Cancer				Other	Total				
		NR	PR	CR	% Response	NR	PR	CR	% Response	NR	PR	CR	% Response	NR	NR	PR	CR	% Response	
3	1	2	0	0	0	3	0	1	25	—	—	—	—	—	5	0	1	17	
3	3	11	2	1	21	9†	2	1	25	2	0	0	0	3§	25	4	2	19	
3	4.5	5	4	0	44	4‡	2	0	33	5	0	0	0	5§	19	6	0	24	
6	4.5	8*	4	2	43	8	3	2	38	1	1	0	50	—	17	8	4	41	
6 (×1)	3	2	2	1	60	6**	4	0	40	1	0	0	0	4†	13	6	1	35	
6 (×1)	6	31	3	3	16	11††	5	1	35	5	0	0	0	5#	51	8	4	19	
Total		59	15	7	27	41	16	5	34	14	1	0	7	17	131	32	12	25	

*Plus one unassessable patient.

†Plus one patient dead of treatment.

‡Plus one unassessable patient.

§One patient each with lymphoma, prostate, and testicular cancer.

||Three patients with breast cancer; one esophageal cancer; one sarcoma.

†One patient each with breast, sarcoma, small bowel, and prostate cancer.

#Two patients with breast cancer; one small bowel; one sarcoma; one Von Hippel Lindau syndrome.

**Plus one patient treated with kidney in place; no response.

††Plus 11 patients treated with kidney in place; one CR, one PR, nine NR.

replicate along with the cell so that daughter cells would have an equal amount of label. Further, this marker could be used to reisolate the infused cells because it would confer resistance to an antibiotic in cells not bearing the gene. In addition, the polymerase chain reaction technique could be used to detect as few as one modified cell in 10,000 normal cells.

Because foreign genes had never been inserted into humans, an extensive review process was undertaken before beginning these gene transfers, including review by the Clinical Research Committees of the NCI, the National Heart, Lung and Blood Institute, the National Institutes of Health Biosafety Committee, the Recombi-

nant DNA Advisory Committee and its Gene Therapy Subcommittee, and the Food and Drug Administration (FDA). A variety of studies were performed and presented to these review groups.^{133,134} We demonstrated that the neomycin phosphotransferase gene could be inserted and expressed in human TIL, that the phenotype and function of marked TIL were not significantly altered, that we could detect the marked gene in animal models, and that there was low risk to the patient and no risk to health care personnel or the public in the use of these techniques. Insertion of the gene into human TIL was assessed by Southern blot analysis. We demonstrated that the gene was expressed and that these transduced cells were resistant to exposure to G418, a neomycin analog normally lethal to all eukaryotic cells (Fig 12). Permission was given to treat 10 patients with life expectancies of up to 90 days.

The first patient received an infusion of gene-modified cells on May 22, 1989. We have now treated 10 patients with advanced melanoma using these gene-modified cells. The results of the first five patients have been published previously.¹³⁵ These patients had extensive melanoma with multiple lesions including metastases in brain, lung, liver, subcutaneous tissue, adrenal gland, and other sites (Table 12). The characteristics of the infused gene-transduced and nontransduced TIL administered are listed in Table 13. The primary goal of these studies was the detection of these gene-marked cells in blood and tumor samples. A summary of the polymerase chain reaction analysis of the distribution of gene-modified TIL in peripheral blood and in tumor in the first five patients is shown in Fig 13. We have been

Table 11. Experimental Leads for Improving Immunotherapy

- Enhancing in vivo effectiveness of TIL
 - Combination with other cytokines
 - Combination with local radiation therapy
- Generating more effective TIL
 - Lymphocyte subpopulations or clones
 - Repeated in vitro stimulation
 - Low-dose IL-2
 - Culture in IL-2 plus IL-4
 - Modify TIL by gene transfer
- Immunize with gene-modified tumor
 - Cytokine genes
 - MHC genes
- Application of new cytokines
 - IL-6
 - Cytokine combinations
- Monoclonal antibodies
 - +IL-2 + LAK cells (LAK cells mediate ADCC)
 - +M-CSF
- Synergy of chemotherapy + IL-2

Abbreviations: ADCC, antibody-dependent cell-mediated cytotoxicity; M-CSF, macrophage colony-stimulating factor.




Fig 12. Transduced TIL become resistant to exposure to G418. Nontransduced TIL grow in the absence (●) but not in the presence (○, 0.4 mg/mL) of G418, a neomycin analog. Transduced TIL grow well in the presence and absence of G418.

able to detect gene-modified cells in peripheral blood at times up to 189 days and in tumor up to 64 days after cell infusion.¹³⁵ In general, however, gene-modified cells are only found in the circulation for the first 3 weeks after infusion, which corresponds to the time that IL-2 is administered. Two of the first five patients had objective

responses to gene-modified TIL. The patient we previously reported had undergone complete regression of multiple subcutaneous, lung, and mucosal melanoma deposits remains alive and free of disease beyond 2 years.

All of the safety studies performed on the retroviral supernatant used for transduction and the transduced cells infused into the patient and multiple tests on patients at varying intervals after the infusion of transduced cells revealed no safety problems of any kind. No patient was ever exposed to replication-competent retrovirus, as evidenced by sensitive S+ L- 3T3 amplification assays or tests for antibody to envelope proteins in patient serum.¹³⁵

These studies demonstrated that retroviral-mediated gene transduction could be safely used to modify genetically cells used for infusion into patients and led to two applications to attempt to improve the therapy of established disease. Because lymphocytes appeared to be suitable vehicles for introducing genes, clinical efforts led by Drs R. Michael Blaese and W. French Anderson were initiated to insert the gene for adenosine deaminase into children with severe combined immunodeficiency, resulting from a deficiency of this enzyme. A second series of efforts were devoted to the gene therapy of cancer. An implication of our studies using gene transduction into TIL was the concept that gene modification might be used to design TIL with new properties that could increase their therapeutic effectiveness. Several possibilities existed for the genetic modification of TIL and some of these are listed in Table 14. The first

Table 12. Patient Characteristics

Patient No.	Age (years)/ Sex	Primary Site	Prior Treatment	Site	Size (cm)	Tumor Harvest				Sites of Assessable Disease
						Total No. of Cells Obtained ($\times 10^{-7}$)	No. of Cells to Start Culture* ($\times 10^{-7}$)	% Lymphocytes	% Tumor Cells	
1	52/M	Neck	Wide local excision	Lymph node	4 × 4 × 2	33	12	15	85	Lung, liver, spleen
2	46/F	Finger	Amputation finger lymph node dissection	Lymph node	5 × 5 × 3	157	50	70	30	Lymph nodes, intramuscular
3	42/M	Back	Wide local excision	Lymph node	6 × 5 × 4	205	120	31	69	Lung, subcutaneous
			Lymph node dissection	Subcutaneous	2 × 2 × 2					
			Melanoma vaccine IL-2/interferon alfa	Subcutaneous	2 × 2 × 2					
4	41/M	Chest	Wide local excision	Subcutaneous	2 × 1 × 1	41	4	39	61	Lung, liver, lymph nodes, subcutaneous, brain
			Lymph node dissection	Subcutaneous	5 × 4 × 4					
			IL-2/interferon alfa							
5	26/F	Arm	Wide local excision	Subcutaneous	2 × 2 × 2 to 5 × 4 × 2	71	15	16	84	Lung, lymph nodes, subcutaneous

NOTE. Reprinted with permission from Rosenberg et al.¹

*Denotes the number of cells used to start the cultures that were ultimately administered to the patient.

Table 13. Characteristics of Infused Cells

Patient No.	Cells Transduced/Day of Transduction	No. of Cells Transduced ($\times 10^{-8}$)	Multiplicity of Infection*	Total Days of Growth	Fold Expansion†	Doubling Time‡ (days)	No. of Cells Infused ($\times 10^{-10}$)	Estimate of Cells Transduced (%)	Cycle Administered	No. of IL-2 Doses to Patient
1	No	—	—	36 and 37	63,400	2.5	22.8	—	1	7
	Yes/13	1.8	2.3	60	16,100	7.5	7.1	1	2	
2	No	—	—	65	209	3.5	0.2	—	1	15
	Yes/19	2.5	1.3	65	5,030	3.5	13.2	11	1	
3§										
A	No	—	—	35	874	2.8	10.8	—	1	13
	Yes/12	1.8	1.6	48	35,100	2.5	14.5	1	2	
B	No	—	—	35	324	2.3	3.5	—	1	
	Yes/12	1.4	1.6	48	21,400	3.5	5.5	1	2	
4	No	—	—	36	8,470	2.0	26.0	—	1	10
	Yes/16	1.8	1.7	36	9,480	2.0	3.3	4	1	
5	No	—	—	30	18,900	2.0	15.0	—	1	7
	Yes/8	0.6	1.6	30	5,250	2.3	6.2	10	1	

NOTE. Reprinted with permission from Rosenberg et al.¹

*Ratio of number of virions to number of TIL during the transduction procedure.

†Calculated fold expansion of cells administered. Not all cultured cells were given; some cells were diverted for experimental studies or lost to contamination. Nontransduced cells shown here are those in conjunction with the transduced cells administered to the patient.

‡Varied during culture growth; doubling time at time of final cell infusion is presented here.

§Two separate cultures transduced and expanded separately. Culture A was started in AIM-V medium; culture B was started in RPMI 1640 plus 10% human serum.

gene we selected was TNF because animal and laboratory experiments suggested that TIL secreting large amounts of TNF might provide therapeutic benefit to patients with established cancer.

TNF is an extremely effective therapeutic agent for




Fig 13. Results of polymerase chain reaction assays of peripheral blood lymphocytes (PBL) and tumor from cancer patients after infusion of transduced TIL. Consistent demonstration of transduced TIL in the circulation was seen for the first 3 weeks after TIL administration, which corresponded to the time of IL-2 administration. (○, □) Negative; (●, ■) positive. Reprinted, by permission of *The New England Journal of Medicine* (323:570-578, 1990).¹³⁵

the treatment of established mouse tumors in the skin and liver.¹³⁶⁻¹³⁹ These tumors can undergo regression within hours after injection of a single dose of TNF. However, extensive studies in humans have failed to reveal any benefit of TNF administration in humans. In our own phase I studies, 39 patients were treated with escalating doses of TNF to maximum tolerance, and no antitumor effects were seen.⁴² There is a substantial difference in the dose of TNF tolerated by mice and humans. Approximately 400 mg/kg are required to mediate tumor regression in mice. However, humans will tolerate only 8 to 10 mg/kg before life-threatening hypotension supervenes. Because TIL accumulate in tumor deposits, we hypothesized that TIL could be used to produce large amounts of TNF at the site of tumor without exposing the entire patient to TNF. Our calculations predicted that in humans we could achieve at the

Table 14. Possible Genetic Modifications of TIL to Improve Antitumor Activity

Tumor necrosis factor (interferes with blood supply)
Interferon alpha or gamma (upregulated MHC antigens)
Other cytokines such as IL-6, IL-1 alpha, IL-7, RANTES (modulates immune response to tumors)
Fc receptor (mediates antibody-dependent cellular cytotoxicity)
Chimeric T-cell receptors (constant region of T-cell receptor plus variable region of monoclonal antibody alters specificity of T cell)
IL-2 receptor (increase sensitivity to administered IL-2)

tumor site, two to three times the 400 µg/kg required to see tumor regression in mice.

With this aim in mind, we selected a retroviral construct in which the TNF gene was promoted by the viral long terminal repeat promoter, and the neomycin resistance gene was promoted by the SV40 early promoter. These retroviral vectors were supplied to us by Genetic Therapy Inc (Rockville, MD) and Dr Michael Kreigler (Cetus Corp, Emeryville, CA). Extensive testing in vitro revealed that TIL transduced with this TNF vector could secrete high levels of TNF. After considerable review, a protocol for administering these TNF gene-modified TIL was agreed upon by the review groups and the FDA. This protocol schema is shown in Table 15.

Increasing doses of TNF-TIL are given without IL-2 starting at very low doses (10^8 cells) and escalated to 10^{11} cells. After these infusions are completed, the dose of TNF/TIL is dropped by a factor of 10 and then administered along with IL-2.

On January 29, 1991, the first two patients with advanced melanoma were treated with TNF gene-modified TIL. The first four patients to receive TNF/TIL are shown in Table 16. Only one of these patients has received gene-modified TIL along with IL-2. No side effects have been seen due to the TNF/TIL. Results are very preliminary, and it will take approximately 2 years to accrue 50 patients to evaluate this therapeutic approach. As of October 1, 1991, one of the patients treated with TNF-TIL alone without IL-2 died of progressive disease 7 months after the last TIL infusion. The only patient to receive TNF-TIL along with IL-2 has shown an objective regression of tumor, which is continuing at the present time.

We are currently exploring a variety of other means for genetically modifying TIL to increase their therapeutic effectiveness (Table 14).

Table 15. Protocol Schema for Administration of TNF-Gene Modified TIL

1. Escalate TNF TIL twice weekly; no IL-2:
10^8 cells
3×10^8
10^9
3×10^9
10^{10}
3×10^{10}
10^{11}
3×10^{11}
2. Reduce cell dose to $\frac{1}{10}$ maximum-tolerated dose. Escalate as in (1) every 3 weeks with 180,000 IU/kg IL-2.
3. After three patients discuss with the FDA (possibly start at a higher cell dose with IL-2).

Table 16. Gene Therapy With TIL Transduced With the Gene for TNF

Patient No.	Age (years)/Sex	Sites of Metastatic Disease	Secretion of TNF by TIL (pg/ 10^6 cells/24h)	Cell Doses Administered
1	30/F	Brain, subcutaneous, adrenal	141	1.1×10^8 3.0×10^8 1.0×10^9 3.0×10^9 5.4×10^9
2	44/M	Lymph nodes, subcutaneous, intramuscular	517	1.0×10^8 3.1×10^8 1.1×10^9 3.0×10^9
3	53/F	Liver, subcutaneous, retroperitoneum	125	1.0×10^8 3.0×10^8 9.6×10^8 3.0×10^9 1.0×10^{10} 2.4×10^{10} 1.0×10^{11} $1.0 \times 10^{10} + \text{IL-2}$ $2.8 \times 10^{10} + \text{IL-2}$ $9.0 \times 10^{10} + \text{IL-2}$
4	25/M	Brain, lung, subcutaneous, mediastinum	115	1.0×10^9 1.0×10^{10}

tic effectiveness (Table 14). The introduction of other cytokines such as interferon alfa or gamma can upregulate MHC antigens and may synergize with TIL activity. The introduction of Fc receptors into TIL may allow their use in conjunction with monoclonal antibodies to mediate antibody-dependent cellular cytotoxicity. A particularly intriguing application of TIL is the introduction of chimeric T-cell receptors to modify the antigenic range of these cells. These chimeric receptors consist of the constant region of the T-cell receptor plus the variable region of the monoclonal antibody. Gross et al¹⁴⁰ have shown that the introduction of these chimeric T-cell receptors into mouse T-cell hybridomas can alter the antigenic specificity of these cells.¹⁴⁰ The use of this technology in conjunction with the use of TIL could potentially increase the range of TIL activity to tumors for which monoclonal antibodies exist, and we are actively exploring this potential application. The introduction of IL-2 receptors into TIL might increase their sensitivity to IL-2 and, thus, enable us to use lower levels of administered IL-2.

More recently we have begun studies of the genetic modification of cancer cells to increase their immunogenicity. A variety of investigators have shown that introduction of cytokine genes into tumor cells can increase

the recognition of tumor by the host immune system.¹⁴¹⁻¹⁴⁸ The cytokine genes used in these studies have included IL-4, IL-2, interferon gamma, TNF, and granulocyte-macrophage colony-stimulating factor. The introduction of genes coding for MHC antigens can also increase tumor immunogenicity. Our work is centered on introducing the genes for TNF and IL-2 into tumor cells. Mouse tumor cells transduced with the TNF gene grow for a short time and then spontaneously regress. This regression is dependent on the presence of CD4+ and CD8+ cells, as has been demonstrated by antibody-depletion studies. We have thus designed a clinical protocol for the treatment of patients with advanced cancer. The protocol schema is outlined in Table 17. Tumor is excised from patients, and tissue culture lines established. Genes for either TNF or IL-2 are inserted into these tumor cells, and these tumor-cell vaccines are then used to immunize the patient against their autologous cancer. In addition to this active immunization, draining lymph nodes will be excised 3 weeks after tumor-cell injection grown in culture and used for adoptive immunotherapy.

The studies described in this report demonstrate that it is possible to manipulate the host immune system to

Table 17. Protocol Schema for the Use of Tumor Cells Transduced With the Genes for TNF or IL-2 for the Treatment of Patients With Advanced Cancer

1. Tumor resected as part of standard treatment.
2. Tissue culture line established.
3. Cytokine gene transduced into culture, and cultures selected in G418 (cytokine > 100 pg/10⁶ cells/24h).
4. Inject 2×10^8 gene-modified tumor cells subcutaneously into right thigh and 2×10^7 cells intradermally at two nearby sites.
5. Three weeks later, remove draining lymph node and grow in vitro in IL-2.
6. Adoptive immunotherapy using these cells plus IL-2.

mediate the regression of cancer in some patients with advanced disease. The immune manipulations we are currently using are in the infancy of their development and as increased knowledge accrues concerning the nature of the immune response to human tumors, more sophisticated approaches should be possible. The fact that at least some bulky human tumors can be impacted by immunologic maneuvers is encouraging for future progress in this field. With future developments, it is hoped that these early approaches can be developed into effective, safe, and practical treatments for patients with cancer.

REFERENCES

1. Rosenberg SA, Longo DL, Lotze MT: Principles and applications of biologic therapy, in DeVita VT, Hellman S, Rosenberg SA (eds): *Cancer, Principles and Practices of Oncology*, ed 3. Philadelphia, PA, Lippincott, 1989, pp 1342-1398
2. DeVita V, Hellman S, Rosenberg SA (eds): *Biologic Therapy of Cancer*. Philadelphia, PA, Lippincott, 1991
3. Rosenberg SA, Terry W: Passive immunotherapy of cancer in animals and man. *Adv Cancer Res* 25:323-388, 1977
4. Morgan DA, Ruscetti FW, Gallo RG: Selective in vitro growth of T-lymphocytes from normal bone marrow. *Science* 193:1007-1008, 1976
5. Smith KA: Interleukin-2: Inception, impact, and implications. *Science* 240:1169-1176, 1988
6. Yron I, Wood TA, Spiess PJ, et al: In vitro growth of murine T cells. V. The isolation and growth of lymphoid cells infiltrating syngeneic solid tumors. *J Immunol* 125:238-245, 1980
7. Lotze MT, Line BR, Mathisen DJ, et al: The in vivo distribution of autologous human and murine lymphoid cells grown in T cell growth factor (TCGF): Implications for the adoptive immunotherapy of tumors. *J Immunol* 125:1487-1493, 1980
8. Grimm EA, Mazumder A, Zhang HZ, et al: Lymphokine-activated killer cell phenomenon: Lysis of natural killer-resistant fresh solid tumor cells by interleukin-2 activated autologous human peripheral blood lymphocytes. *J Exp Med* 155:1823-1841, 1982
9. Grimm EA, Ramsey KM, Mazumder A, et al: Lymphokine-activated killer cell phenomenon. II. Precursor phenotype is serologically distinct from peripheral T lymphocytes, memory cytotoxic thymus-derived lymphocytes, and natural killer cells. *J Exp Med* 157:884-897, 1983
10. Grimm EA, Robb RJ, Roth JA, et al: Lymphokine-activated killer cell phenomenon. III. Evidence that IL-2 is sufficient for direct activation of peripheral blood lymphocytes into lymphokine-activated killer cells. *J Exp Med* 158:1356-1361, 1983
11. Rosenstein M, Yron I, Kaufmann Y, et al: Lymphokine-activated killer cells: Lysis of fresh syngeneic natural killer-resistant murine tumor cells by lymphocytes cultured in interleukin-2. *Cancer Res* 44:1946-1953, 1984
12. Rayner AA, Grimm EA, Lotze MT, et al: Lymphokine-activated killer (LAK) cells: Analysis of factors relevant to the immunotherapy of human cancer. *Cancer* 55:1327-1333, 1985
13. Rosenberg SA: Adoptive immunotherapy of cancer using lymphokine activated killer cells and recombinant interleukin-2, in DeVita VT, Hellman S, Rosenberg SA (eds): *Important Advances in Oncology 1986*. Philadelphia, PA, Lippincott, 1986, pp 55-91
14. Rayner AA, Grimm EA, Lotze MT, et al: Lymphokine activated killer (LAK) cell phenomenon. IV. Lysis by LAK cell clones of fresh human tumor cells from autologous and multiple allogeneic tumors. *J Natl Cancer Inst* 75:67-75, 1985
15. Yang JC, Mulé JJ, Rosenberg SA: Murine lymphokine-activated killer cells. *J Immunol* 137:715-722, 1985
16. Roberts K, Lotze MT, Rosenberg SA: Separation and functional studies of the human lymphokine-activated killer cell. *Cancer Res* 47:4366-4371, 1987
17. Skibber JM, Lotze MT, Uppenkamp I, et al: Identification and expansion of human lymphokine-activated killer cells: Implications for the immunotherapy of cancer. *J Surg Res* 42:613-621, 1987

18. Fox B, Rosenberg SA: Heterogeneous lymphokine-activated killer cell precursor populations. *Cancer Immunol Immunother* 29:155-166, 1989
19. Mazumder A, Rosenberg SA: Successful immunotherapy of NK-resistant established pulmonary melanoma metastases by the intravenous adoptive transfer of syngeneic lymphocytes activated in vitro by interleukin-2. *J Exp Med* 159:495-507, 1984
20. Mulé JJ, Shu S, Schwarz SL, et al: Adoptive immunotherapy of established pulmonary metastases with LAK cells and recombinant interleukin-2. *Science* 225:1487-1489, 1984
21. Mulé JJ, Shu S, Rosenberg SA: The anti-tumor efficacy of lymphokine-activated killer cells and recombinant interleukin-2 in vivo. *J Immunol* 135:646-652, 1985
22. Rosenberg SA, Mulé JJ, Spiess PJ, et al: Regression of established pulmonary metastases and subcutaneous tumor mediated by the systemic administration of high dose recombinant IL-2. *J Exp Med* 161:1169-1188, 1985
23. Lafreniere R, Rosenberg SA: Successful immunotherapy of murine experimental hepatic metastases with lymphokine-activated killer cells and recombinant interleukin-2. *Cancer Res* 45:3735-3741, 1985
24. Lafreniere R, Rosenberg SA: Adoptive immunotherapy of murine hepatic metastases with lymphokine activated killer (LAK) cells and recombinant interleukin-2 (RIL-2) can mediate the regression of both immunogenic and non-immunogenic sarcomas and an adenocarcinoma. *J Immunol* 135:4273-4280, 1985
25. Mulé JJ, Ettinghausen SE, Spiess PJ, et al: The anti-tumor efficacy of lymphokine-activated killer cells and recombinant interleukin-2 in vivo: An analysis of survival benefit and mechanisms of tumor escape in mice undergoing immunotherapy. *Cancer Res* 46:676-683, 1986
26. Mulé JJ, Yang J, Shu S, et al: The anti-tumor efficacy of lymphokine-activated killer cells and recombinant interleukin-2 in vivo: Direct correlation between reduction of established metastases and cytolytic activity of lymphokine-activated killer cells. *J Immunol* 136:3899-3909, 1986
27. Papa MZ, Mulé JJ, Rosenberg SA: The anti-tumor efficacy of lymphokine-activated killer cells and recombinant interleukin-2, in Lapis K, Liotta LA, Rabson AS (ed): *The Biochemistry and Molecular Genetics of Cancer Metastases*. Boston, MA, Martinus Nijhoff, 1986, pp 307-319
28. Papa MZ, Vetto JT, Ettinghausen SE, et al: Effects of corticosteroid on the anti-tumor activity of lymphokine activated killer cells and interleukin-2 in mice. *Cancer Res* 46:5618-5623, 1986
29. Shiloni E, Lafreniere R, Mulé JJ, et al: Effect of immunotherapy with allogeneic lymphokine activated killer (LAK) cells and recombinant interleukin-2 (RIL-2) on established pulmonary and hepatic metastases in mice. *Cancer Res* 46:5633-5640, 1986
30. Ettinghausen SE, Lipford EH, Mulé JJ, et al: Systemic administration of recombinant interleukin-2 stimulates in vivo lymphoid cell proliferation in tissues. *J Immunol* 135:1488-1497, 1985
31. Ettinghausen SE, Lipford EH, Mulé JJ, et al: Recombinant interleukin-2 stimulates in vivo proliferation of adoptively transferred lymphokine activated killer (LAK) cells. *J Immunol* 135:3623-3635, 1985
32. Taniguchi T, Matsui H, Fujita T, et al: Structure and expression of a cloned cDNA for human interleukin-2. *Nature* 302:305-307, 1983
33. Fujita T, Takaoka C, Matsui H, et al: Structure of the human interleukin-2 gene. *Proc Natl Acad Sci USA* 80:7437-7441, 1983
34. Rosenberg SA, Grimm EA, McGroffan M, et al: Biological activity of recombinant human interleukin-2 produced in *Escherichia coli*. *Science* 223:1412-1415, 1984
35. Wang A, Lu S-D, Mark DF: Site-specific mutagenesis of the human interleukin-2 gene: Structure-function analysis of the cysteine residues. *Science* 224:1431-1433, 1984
36. Lotze MT, Frana LW, Sharrow SO, et al: In vivo administration of purified human interleukin-2. I. Half-life and immunologic effects of the Jurkat cell line-derived interleukin-2. *J Immunol* 134:157-166, 1985
37. Lotze MT, Matory YL, Ettinghausen SE, et al: In vivo administration of purified human interleukin-2. II. Half-life, immunologic effects, and expansion of peripheral lymphoid cells in vivo with recombinant IL-2. *J Immunol* 135:2865-2875, 1985
38. Rosenberg SA: Immunotherapy of cancer by the systemic administration of lymphoid cells plus interleukin-2. *J Biol Response Mod* 3:501-511, 1984
39. Rosenberg SA, Lotze MT, Muul LM, et al: Observations on the systemic administration of autologous lymphokine-activated killer cells and recombinant interleukin-2 to patients with metastatic cancer. *N Engl J Med* 313:1485-1492, 1985
40. Lotze MT, Chang AE, Seipp CA, et al: High-dose recombinant interleukin-2 in the treatment of patients with disseminated cancer. *JAMA* 256:3117-3124, 1986
41. Rosenberg SA, Lotze MT, Muul LM, et al: A progress report on the treatment of 157 patients with advanced cancer using lymphokine activated killer cells and interleukin-2 or high dose interleukin-2 alone. *N Engl J Med* 316:889-905, 1987
42. Rosenberg SA, Lotze MT, Yang JC, et al: Experience with the use of high-dose interleukin-2 in the treatment of 652 cancer patients. *Ann Surg* 210:474-485, 1989
43. Rosenstein M, Ettinghausen SE, Rosenberg SA: Extravasation of intravascular fluid mediated by the systemic administration of recombinant interleukin-2. *J Immunol* 137:1735-1742, 1986
44. Lotze MT, Matory YL, Rayner AA, et al: Clinical effects and toxicity of interleukin-2 in patients with cancer. *Cancer* 58:2764-2772, 1986
45. Ettinghausen SE, Moore JG, White DE, et al: Hematologic effects of immunotherapy with lymphokine activated killer cells and recombinant interleukin-2 in cancer patients. *Blood* 69:1654-1660, 1987
46. Vetto JT, Papa MZ, Lotze MT, et al: Reduction of toxicity of interleukin-2 and lymphokine activated killer cells in humans by the administration of corticosteroids. *J Clin Oncol* 5:496-503, 1987
47. Beldegrun A, Webb DE, Austin HA, et al: Effects of interleukin-2 on renal function in patients receiving immunotherapy for advanced cancer. *Ann Intern Med* 106:817-822, 1987
48. Denicoff KD, Rubinow DR, Papa MZ, et al: The neuropsychiatric effects of the treatment with interleukin-2 and lymphokine activated killer cells. *Ann Intern Med* 107:293-300, 1987
49. Cotran RS, Pober JS, Gimbrone MS, Jr, et al: Endothelial activation during interleukin-2 (IL-2) immunotherapy: A possible mechanism for the vascular leak syndrome. *J Immunol* 140:1883-1888, 1988
50. Gaspari AA, Lotze MT, Rosenberg SA, et al: Dermatologic changes associated with interleukin-2 administration. *JAMA* 258:1624-1629, 1987

51. Saris SC, Rosenberg SA, Friedman RB, et al: Penetration of recombinant interleukin-2 across the blood cerebrospinal fluid barrier. *J Neurosurg* 69:29-34, 1988
52. Ognibene FP, Rosenberg SA, Lotze M, et al: Interleukin-2 administration causes reversible hemodynamic changes and left ventricular dysfunction similar to those seen in septic shock. *Chest* 94:750-754, 1988
53. Webb DE, Austin HA, Beldegrun A, et al: Metabolic and renal effects of interleukin-2 immunotherapy for metastatic cancer. *Clin Nephrol* 30:141-145, 1988
54. Schwartzentruber D, Lotze MT, Rosenberg SA: Colonic perforation: An unusual complication of therapy with high-dose interleukin-2. *Cancer* 62:2350-2353, 1988
55. Lee RE, Gaspari AA, Lotze MT, et al: Interleukin-2 and psoriasis. *Arch Dermatol* 124:1811-1815, 1988
56. Lee RE, Lotze MT, Skibber JM, et al: Cardiorespiratory effects of immunotherapy with interleukin-2. *J Clin Oncol* 7:7-20, 1989
57. Beldegrun A, Webb DE, Austin HA, et al: Renal toxicity of interleukin-2 administration in patients with metastatic renal cell cancers: Effect of pretherapy nephrectomy. *J Urol* 141:499-503, 1989
58. Denicoff KD, Durkin TM, Lotze MT, et al: The neuroendocrine effects of interleukin-2 treatment. *J Clin Endocrinol Metab* 69:402-410, 1989
59. Kragel AH, Travis WD, Steis RG, et al: Myocarditis or acute myocardial infarction associated with interleukin-2 therapy for cancer. *Cancer* 66:1513-1516, 1990
60. Bock SN, Lee RE, Fisher B, et al: A prospective randomized trial evaluating prophylactic antibiotics to prevent triple-lumen catheter-related sepsis in patients treated with immunotherapy. *J Clin Oncol* 8:161-169, 1990
61. Atkins MB, Mier JW, Parkinson DR, et al: Hypothyroidism after treatment with interleukin-2 and lymphokine-activated killer cells. *N Engl J Med* 318:1557-1563, 1988
62. Schwartzentruber DJ, White DE, Zweig MH, et al: Thyroid dysfunction associated with immunotherapy for patients with cancer. *Cancer* 68:2384-2390, 1991
63. Hawkins MJ: IL-2/LAK. Principles and Practices of Oncology Updates 3:1-14, 1989
64. Sosman JA, Kohler PC, Hank JA, et al: Repetitive weekly cycles of interleukin-2. II. Clinical and immunological effects of dose, schedule and addition of indomethacin. *J Natl Cancer Inst* 80:1451-1461, 1988
65. Allison MAK, Jones SE, McGuffey P: Phase II trial of outpatient interleukin-2 in malignant lymphoma, chronic lymphocytic leukemia, and selected solid tumors. *J Clin Oncol* 7:75-80, 1989
66. Hersh EM, Murray JL, Hong WK, et al: Phase I study of cancer therapy with recombinant interleukin-2 administered by intravenous bolus injection. *J Clin Oncol* (in press)
67. Creekmore SP, Harris JE, Ellis TM, et al: A phase I clinical trial of recombinant interleukin-2 by periodic 24-hour intravenous infusions. *J Clin Oncol* 7:276-284, 1989
68. Bukowski RM, Goodman P, Crawford ED, et al: Phase II trial of high-dose intermittent interleukin-2 in metastatic renal cell carcinoma: A Southwest Oncology Group study. *J Natl Cancer Inst* 82:143-146, 1990
69. Schoof DD, Gramolini BA, Davidson DL, et al: Adoptive immunotherapy of human cancer using low-dose recombinant interleukin-2 and lymphokine-activated killer cells. *Cancer Res* 48:5007-5010, 1988
70. Dutcher JP, Creekmore S, Weiss GR, et al: A phase II study of interleukin-2 and lymphokine-activated killer cells in patients with metastatic malignant melanoma. *J Clin Oncol* 7:477-485, 1989
71. Thompson JA, Lee DJ, Lindgren CG, et al: Influence of schedule of interleukin-2 administration on therapy with interleukin-2 and lymphokine activated killer cells. *Cancer Res* 49:235-240, 1989
72. West WH, Tauer KW, Yannelli JR, et al: Constant-infusion recombinant interleukin-2 in adoptive immunotherapy of advanced cancer. *N Engl J Med* 316:898-905, 1987
73. Paciucci PA, Holland JF, Glidewell O, et al: Recombinant interleukin-2 continuous infusion and adoptive transfer of recombinant interleukin-2 activated cells in patients with advanced cancer. *J Clin Oncol* 7:869-878, 1989
74. Negrier S, Phillip T, Stoter G, et al: Interleukin-2 with or without LAK cells in metastatic renal cell carcinoma: A report of a European multicentre study. *Eur J Cancer Clin Oncol* 25:S21-S28, 1989
75. Stahel RA, Sculier J, Jost LM, et al: Tolerance and effectiveness of recombinant interleukin-2 (r-met Hu IL-2 (ala 125) and lymphokine-activated killer cells in patients with metastatic solid tumors. *Eur J Cancer Clin Oncol* 25:965-972, 1989
76. Dutcher JP, Gaynor ER, Boldt DH, et al: A phase II study of high-dose continuous infusion interleukin-2 with lymphokine-activated killer cells in patients with metastatic melanoma. *J Clin Oncol* 9:641-648, 1991
77. Rosenberg SA, Spiess P, Lafreniere R: A new approach to the adoptive immunotherapy of cancer with tumor infiltrating lymphocytes. *Science* 233:1318-1321, 1986
78. Spiess PJ, Yang JC, Rosenberg SA: In vivo antitumor activity of tumor-infiltrating lymphocytes expanded in recombinant interleukin-2. *J Natl Cancer Inst* 79:1067-1075, 1987
79. Barth RJ Jr, Bock SN, Mulé JJ, et al: Unique murine tumor associated antigens identified by tumor infiltrating lymphocytes. *J Immunol* 144:1531-1537, 1990
80. Barth RJ, Mulé JJ, Spiess PJ, et al: Interferon gamma and tumor necrosis factor have a role in tumor regressions mediated by murine CD8+ tumor infiltrating lymphocytes. *J Exp Med* 173:647-658, 1991
81. Barth RJ, Mulé JJ, Asher AL, et al: Identification of unique murine tumor associated antigens by tumor infiltrating lymphocytes using tumor specific secretion of interferon-gamma and tumor necrosis factor. *J Immunol Methods* 140:269-279, 1991
82. Rosenberg SA, Schwarz S, Spiess P: Combination immunotherapy of cancer: Synergistic anti-tumor interactions of interleukin-2, alpha-interferon and tumor infiltrating lymphocytes. *J Natl Cancer Inst* 80:1393-1397, 1988
83. Cameron RB, Spiess PJ, Rosenberg SA: Synergistic antitumor activity of tumor infiltrating lymphocytes, interleukin-2 and local tumor irradiation: Studies on the mechanism of action. *J Exp Med* 171:249-263, 1990
84. Yang JC, Perry-Lalley D, Rosenberg SA: An improved method for growing murine tumor infiltrating lymphocytes with in vivo antitumor activity. *J Biol Response Mod* 9:149-159, 1990
85. Alexander RB, Rosenberg SA: Long term survival of adoptively-transferred tumor infiltrating lymphocytes in mice. *J Immunol* 145:1615-1620, 1990

86. Weber JS, Rosenberg SA: Effects of murine tumor I major histocompatibility complex expression on antitumor activity of tumor-infiltrating lymphocytes. *J Natl Cancer Inst* 82:755-761, 1990
87. Wong RA, Alexander RB, Puri RK, et al: In vivo proliferation of adoptively transferred tumor-infiltrating lymphocytes in mice. *J Immunol* 100:120-130, 1991
88. Karpati RM, Banks SM, Malissen B, et al: Phenotypic characterization of murine tumor-infiltrating T lymphocytes. *J Immunol* 146:2043-2051, 1991
89. Alexander RB, Rosenberg SA: Adoptively transferred tumor infiltrating lymphocytes can cure established metastatic tumor in mice and persistent long term in vivo as functional memory T lymphocytes. *J Immunother* 10:389-397, 1991
90. Muul LM, Spiess PJ, Director EP, et al: Identification of specific cytolytic immune responses against autologous tumor in humans bearing malignant melanoma. *J Immunol* 138:989-995, 1987
91. Itoh K, Tilden AB, Balch CM: Interleukin-2 activation of cytotoxic T-lymphocytes infiltrating into human metastatic melanomas. *Cancer Res* 46:3011-3017, 1986
92. Topalian SL, Solomon D, Rosenberg SA: Tumor-specific cytotoxicity by lymphocytes infiltrating human melanomas. *J Immunol* 142:3714-3725, 1989
93. Topalian SL, Rosenberg SA: Tumor-infiltrating lymphocytes: Evidence for specific immune reactions against growing cancers in mice and humans, in DeVita VT, Hellman S, Rosenberg SA (eds): *Important Advances in Oncology*. Philadelphia, PA, Lippincott, 1990, pp 19-41
94. Kradin RL, Lazarus DS, Dubinett SM, et al: Tumor-infiltrating lymphocytes and interleukin-2 in treatment of advanced cancer. *Lancet* 1:577-580, 1989
95. Kurnick JT, Kradin RL, Blumberg R, et al: Functional characterization of T lymphocytes propagated from human lung carcinoma. *Clin Immunol Immunopathol* 38:367-380, 1986
96. Rabinowich H, Cohen R, Bruderman I, et al: Functional analysis of mononuclear cells infiltrating into tumors: Lysis of autologous human tumor cells by cultured infiltrating lymphocytes. *Cancer Res* 47:173-177, 1987
97. Galili U, Vanky F, Rodriguez L, et al: Activated T lymphocytes within human solid tumors. *Cancer Immunol Immunother* 6:129-133, 1979
98. Kuppner MC, Hamou MF, de Tribolet N: Immunohistological and functional analyses of lymphoid infiltrates in human glioblastomas. *Cancer Res* 48:6926-6932, 1988
99. Saito T, Tanaka R, Yoshida S, et al: Immunohistochemical analysis of tumor-infiltrating lymphocytes and major histocompatibility antigens in human gliomas and metastatic brain tumors. *Surg Neurol* 29:435-442, 1988
100. Paine JT, Handa H, Yamasaki T, et al: Immunohistochemical analysis of infiltrating lymphocytes in central nervous system tumors. *Neurosurgery* 18:766-772, 1986
101. Heo DS, Whiteside TL, Kanbour A, et al: Lymphocytes infiltrating human ovarian tumors. I. Role of Leu-19 (NKH1)-positive recombinant IL-2 activated cultures of lymphocytes infiltrating human ovarian tumors. *J Immunol* 140:4042-4049, 1988
102. Tagaki S, Chen K, Schwarz R, et al: Functional and phenotypic analysis of tumor-infiltrating lymphocytes isolated from human primary and metastatic liver tumors and cultured in recombinant interleukin-2. *Cancer* 63:102-111, 1989
103. Heo DL, Whiteside TL, Johnson JT, et al: Long-term interleukin-2-dependent growth and cytotoxic activity of tumor-infiltrating lymphocytes from human squamous cell carcinomas of the head and neck. *Cancer Res* 47:6353-6362, 1987
104. Whiteside TL, Miescher S, Hurlimann J, et al: Separation, phenotyping, and limiting dilution analysis of T-lymphocytes infiltrating human solid tumors. *Int J Cancer* 37:803-811, 1986
105. Whiteside TL, Miescher S, Hurlimann J, et al: Clonal analysis and in situ characterization of lymphocytes infiltrating human breast carcinomas. *Cancer Immunol Immunother* 23:169-178, 1986
106. Radrizzani M, Gambacorti-Passerini C, Parmiani G, et al: Lysis by interleukin-2-stimulated tumor-infiltrating lymphocytes of autologous and allogeneic tumor target cells. *Cancer Immunol Immunother* 28:67-73, 1989
107. Itoh K, Platsoucas CD, Balch CM: Autologous tumor-specific cytotoxic T lymphocytes in the infiltrate of human metastatic melanomas: Activation by interleukin 2 and autologous tumor cells and involvement of the T cell receptor. *J Exp Med* 168:1419-1441, 1988
108. Schwartzentruber DJ, Topalian SL, Mancini MJ, et al: Specific release of granulocyte-macrophage colony-stimulating factor, tumor necrosis factor- α , and IFN- γ by human tumor-infiltrating lymphocytes after autologous tumor stimulation. *J Immunol* 146:153-164, 1991
109. Darrow TL, Slingluff CL, Seigler HF: The role of class I antigens in recognition of melanoma cells by tumor-specific cytotoxic T lymphocytes: Evidence for shared tumor antigens. *J Immunol* 142:3329-3335, 1989
110. Hom SS, Topalian SL, Simoni ST, et al: Common expression of melanoma tumor-associated antigens recognized by human tumor-infiltrating lymphocytes: Analysis by HLA restriction. *J Immunother* 10:153-164, 1991
111. Topalian SL, Solomon D, Avis FP, et al: Immunotherapy of patients with advanced cancer using tumor-infiltrating lymphocytes and recombinant interleukin-2: A pilot study. *J Clin Oncol* 6:839-853, 1988
112. Berendt MJ, North RJ: T-cell-mediated suppression of anti-tumor immunity: An explanation for progressive growth of an immunogenic tumor. *J Exp Med* 151:69-80, 1980
113. Greenberg PD, Cheever MA: Treatment of disseminated leukemia with cyclophosphamide and immune cells: Tumor immunity reflects long-term persistence of tumor-specific donor T cells. *J Immunol* 133:3401-3407, 1984
114. Rosenberg SA, Packard BS, Aebbersold PM, et al: Use of tumor-infiltrating lymphocytes and interleukin-2 in the immunotherapy of patients with metastatic melanoma, special report. *N Engl J Med* 319:1676-1680, 1988
115. Fisher B, Packard BS, Read EJ, et al: Tumor localization of adoptively transferred indium-111 labeled tumor infiltrating lymphocytes in patients with metastatic melanoma. *J Clin Oncol* 7:250-261, 1989
116. Griffith KD, Read EJ, Carrasquillo CS, et al: In vivo distribution of adoptively transferred indium-111 labeled tumor infiltrating lymphocytes and peripheral blood lymphocytes in patients with metastatic melanoma. *J Natl Cancer Inst* 81:1709-1717, 1989
117. Rubin JT, Elwood LJ, Rosenberg SA, et al: Immunohistochemical correlates of response to recombinant interleukin-2 based immunotherapy in humans. *Cancer Res* 49:7086-7092, 1989
118. Cameron RB, McIntosh JK, Rosenberg SA: Synergistic antitumor effects of combination immunotherapy with recombinant interleukin-2 and a recombinant hybrid interferon- α in

the treatment of established murine hepatic metastases. *Cancer Res* 48:5810-5817, 1988

119. Rosenberg SA, Lotze MT, Yang JC, et al: Combination therapy with interleukin-2 and alpha-interferon for the treatment of patients with advanced cancer. *J Clin Oncol* 7:1863-1874, 1989

120. Kawakami Y, Rosenberg SA, Lotze MT: Interleukin-4 promotes the growth of tumor-infiltrating lymphocytes cytotoxic for human autologous melanoma. *J Exp Med* 168:2183-2191, 1988

121. Mulé JJ, McIntosh JK, Jablons DM, et al: Antitumor activity of recombinant interleukin-6 in mice. *J Exp Med* 171:629-636, 1990

122. Jicha DL, Schwarz S, Mulé JJ, et al: Interleukin-7 mediates the generation and expansion of murine allosensitized and antitumor CTL. *J Immunol* (in press)

123. Eisenthal A, Rosenberg SA: Characterization of IL-2 induced murine cells which exhibit ADCC activity. *Cell Immunol* 115:257-272, 1988

124. Eisenthal A, Lafreniere R, Lefor AT, et al: Effect of anti-B16 melanoma monoclonal antibody on established murine B16 melanoma liver metastases. *Cancer Res* 47:2771-2776, 1987

125. Eisenthal A, Cameron RB, Uppenkamp I, et al: Effect of combined therapy with lymphokine-activated killer cells, interleukin-2 and specific monoclonal antibody on established B16 melanoma lung metastases. *Cancer Res* 48:7140-7145, 1988

126. Papa MZ, Yang JC, Vetto JT, et al: Combined effects of chemotherapy and interleukin-2 in the therapy of mice with advanced pulmonary tumors. *Cancer Res* 48:122-129, 1988

127. Rosenberg SA: Gene therapy of cancer, in DeVita VT, Hellman S, Rosenberg SA: Important Advances in Oncology. Philadelphia, PA, Lippincott (in press)

128. Kriegler M: Gene Transfer and Expression. A Laboratory Manual. New York, NY, Stockton Press, 1990, pp 1-242

129. Gilboa E, Eglitis MA, Kantoff PW, et al: Transfer and expression of cloned genes using retroviral vectors. *BioTechniques* 4:504-512, 1986

130. Eglitis MA, Anderson WF: Retroviral vectors for introduction of genes into mammalian cells. *BioTechniques* 6:608-614, 1988

131. Miller AD, Buttimore C: Redesign of retrovirus packaging cell lines to avoid recombination leading to helper virus production. *Mol Cell Biol* 6:2895-2902, 1986

132. Miller AD, Rosman GJ: Improved retroviral vectors for gene transfer and expression. *BioTechniques* 7:980-986, 1989

133. Kasid A, Morecki S, Aebersold P, et al: Human gene transfer: Characterization of human tumor-infiltrating lymphocytes as vehicles for retroviral-mediated gene transfer in man. *Proc Natl Acad Sci USA* 87:473-477, 1990

134. Culver K, Cornetta K, Morgan R, et al: Lymphocytes as cellular vehicles for gene therapy in mouse and man. *Proc Natl Acad Sci USA* 88:3155-3159, 1991

135. Rosenberg SA, Aebersold P, Cornetta K, et al: Gene transfer into humans—Immunotherapy of patients with advanced melanoma, using tumor-infiltrating lymphocytes modified by retroviral gene transduction. *N Engl J Med* 323:570-578, 1990

136. Carswell EA, Old LJ, Kassel RC, et al: An endotoxin-induced serum factor that causes necrosis of tumors. *Proc Natl Acad Sci USA* 72:3666-3670, 1975

137. Wang AM, Creasy AA, Ladner MB, et al: Molecular cloning of the complementary DNA for human tumor necrosis factor. *Science* 228:149-154, 1985

138. Asher AL, Mulé JJ, Reichert CM, et al: Studies of the anti-tumor efficacy of systemically administered recombinant tumor necrosis factor against several murine tumors in vivo. *J Immunol* 138:963-974, 1987

139. McIntosh JE, Mulé JJ, Merino MJ, et al: Synergistic antitumor effects of immunotherapy with recombinant interleukin-2 and recombinant tumor necrosis factor-alpha. *Cancer Res* 48:4011-4017, 1988

140. Gross G, Waks T, Eshhar Z: Expression of immunoglobulin-T-cell receptor chimeric molecules as functional receptors with antibody-type specificity. *Proc Natl Acad Sci USA* 86:10023-10028, 1989

141. Tepper RI, Pattengale PK, Leder P: Murine interleukin-4 displays potent anti-tumor activity in vivo. *Cell* 57:503-512, 1989

142. Fearon ER, Pardoll DM, Itaya T, et al: Interleukin-2 production by tumor cells bypasses T helper function in the generation of an antitumor response. *Cell* 60:397-403, 1990

143. Gansbacher B, Zier K, Daniels B, et al: Interleukin-2 gene transfer into tumor cells abrogates tumorigenicity and induces protective immunity. *J Exp Med* 172:1217-1224, 1990

144. Ley V, Roth C, Langlade-Demoyen P, et al: A novel approach to the induction of specific cytolytic T cells in vivo. *Res Immunol* 141:855-863, 1990

145. Blankenstein T, Qin Z, Uberla K, et al: Tumor suppression after tumor cell-targeted tumor necrosis factor alpha gene transfer. *J Exp Med* 173:1047-1052, 1991

146. Teng MN, Park BH, Koeppe HKW, et al: Long-term inhibition of tumor growth by tumor necrosis factor in the absence of cachexia or T-cell immunity. *Proc Natl Acad Sci USA* 88:3535-3539, 1991

147. Gansbacher B, Bannerji R, Daniels B, et al: Retroviral vector-mediated γ -interferon gene transfer into tumor cells generates potent and long lasting antitumor immunity. *Cancer Res* 50:7820-7825, 1990

148. Colombo MP, Gerrari G, Stoppacciaro A, et al: Granulocyte colony-stimulating factor gene transfer suppresses tumorigenicity of a murine adenocarcinoma in vivo. *J Exp Med* 173:889-897, 1991