



- [About Us](#)
- [Editorial Board](#)
- [Current Issue](#)
- [Past Issues](#)
- [For Authors](#)
- [Submission](#)
- [Privacy and Confidentiality](#)

## Search

## User Login

☐ Remember Me

- [Lost your password](#)



# Recent Advances in Renal Cell Carcinoma Associated with Xp11.12 Translocations/TFE Gene Fusions

January 25, 2012

[*N A J Med Sci.* 2012;5(1):43-47.] [PDF File](#)

Yunguang Liu, MD, PhD; Bo Xu, MD, PhD; Frank Chen, MD, PhD\*

**Renal cell carcinoma associated with Xp11.2 translocations/TFE gene fusions (Xp11.2 RCC) has been classified as a distinct entity in the 2004 WHO classification of kidney tumors. Over the past seven years, aided by increased awareness, positive nuclear staining for TFE3, unique cytogenetic features, and modern molecular technology, more cases have been recognized as Xp11.2 RCC. This review summarizes the most recent advances in Xp11.2 RCC regarding to its clinical presentation, cytogenetic profile, histopathology, prognosis and treatment.**

**Key Words:** *renal cell carcinoma, Xp11.2 translocations/TFE gene fusions, renal cell carcinoma, Xp11.2 translocations/TFE gene fusions, clinical presentation, cytogenetic profile, histopathology, prognosis and treatment*

Go To: | [Top](#) | [Abstract](#) | [Text](#) | [Author Information](#) | [References](#) |

## INTRODUCTION

Renal cell carcinoma associated with Xp11.2 translocations and TFE3 fusions (Xp11.2 RCC) was first reported by de Jong et al. in 1986.<sup>1</sup> However, it was only recognized as a distinct entity in 2004 WHO classification of kidney tumors.<sup>2</sup> These renal carcinomas are defined by several different translocations involving chromosome Xp11.2 and resulting in gene fusion/overexpression of transcription factor E3 (TFE3) gene. TFE3 is located on chromosome Xp11.2 and belongs to the microphthalmia transcription factor (MiTF)/transcription factor E (TFE) family which also includes 3 other members: MiTF, transcription factor EB (TFEB) and transcription factor EC (TFEC).<sup>3-6</sup> The MiTF/TFE family encodes basic helix-loop-helix-leucine zipper (bHLH-ZIP) transcriptional factors. In addition to regulating melanocytic differentiation, these transcriptional factors also play an important role in proliferation and survival.<sup>7</sup> Normally, the expression of TFE3 is tightly controlled and routine immunohistochemical staining is not able to detect it. A common feature of Xp11.2 RCC is chromosome translocations resulting TFE3 fusion with various partners and subsequently overexpression of TFE3 protein in tumor cells. The overexpressed TFE3 protein in Xp11.2 RCC is now detectable using a sensitive (97.5%) and specific (99.6%) polyclonal antibody to its C-terminal.

## CLINICAL PRESENTATION

The profile of pediatric renal neoplasm is different from that of adult with RCC accounting for <3% of pediatric renal tumors<sup>9-10</sup> and >90% of adult renal tumors.<sup>2</sup> Xp11.2 RCC is estimated representing one-third of pediatric RCC<sup>11</sup> and <1% of adult RCC,<sup>12</sup> and affects patients ranging from 17-month-old<sup>13</sup> to 78-year-old with unknown gender preference in pediatric age group and a female predominance in adults.<sup>14-17</sup> The clinical presentation of Xp11.2 RCC is similar to other renal tumors. Pediatric and young adult patients are usually symptomatic at presentation and only a few cases are incidentally discovered during abdominal imaging. The most common symptom is hematuria, followed by abdominal mass, abdominal pain and weight loss. Rare atypical presentations in adult patients include a heavily calcified renal mass, outflow obstruction with persistent pyelonephritis, renal cyst or nephrolithiasis.<sup>2,15-18</sup> The radiological findings of Xp11.2 RCC are not specific either.<sup>19</sup> However, in young patients, Xp11.2 RCC should be suspected if prominent lymph node metastases are present or the imaging findings are similar to those of papillary RCC.<sup>20-22</sup> In children, studies have indicated that previous chemotherapy may be a risk factor for developing Xp11.2 RCC. Approximately 10-15% of pediatric cases have a history of chemotherapy.<sup>23-25</sup> In adults, even though Xp11.2 RCC has been reported during pregnancy<sup>26</sup> or in association with hemodialysis,<sup>27</sup> no studies have been done to identify particular risk factors.

## CYTOGENETIC PROFILE

To date, eight TFE3 fusion partners have been reported, including papillary renal cell carcinoma (PRCC),<sup>15</sup> alveolar soft part sarcoma locus (ASPL),<sup>16,28</sup> polypyrimidine tract-binding protein-associated splicing factor (PSF),<sup>15</sup> non-POU domain-containing octamer-binding (NonO),<sup>15</sup> clathrin heavy chain (CLTC)<sup>15,29</sup> and three unknown genes.<sup>15,26,30</sup> The most common translocations are ASPL-TFE3, PRCC-TFE3 and PSF-TFE3 fusions (**Table 1**).<sup>18,24-25,31</sup> Two other neoplasms that bear similar translocations are worth mentioning here. One is alveolar soft part sarcoma (ASPS). Both ASPL-TFE3 RCC and ASPS harbor the same ASPL-TFE3 fusion gene. However, the translocation is balanced in ASPL-TFE3 RCC and is unbalanced in ASPS.<sup>32</sup> The other one is t(6;11)(p21;q12) translocated RCC [t(6;11) RCC] that bears Alpha-TFEB fusion gene with overexpression of TFEB. Since TFE3 and TFEB are closely related members of MiTF/TFE transcriptional family, the clinical and histomorphologic features of t(6;11) RCC and Xp11.2 RCC overlap.<sup>18</sup> As a common transcriptional target of TFE3 and TFEB, Cathepsin-K is overexpressed in both t(6;11) RCC and Xp11.2 RCC, but not in other types of RCCs.<sup>33</sup> The immunohistochemical character in distinguishing these two entities is positive nuclear staining for TFE3 in Xp11.2 RCC and for TFEB in t(6;11) RCC.<sup>11,34</sup> The major role of chimeric TFE3 fusion proteins in Xp11.2 RCC is transcriptional dysregulation. Tsuda et al.<sup>35</sup> reported that ASPL-TFE3, PSF-TFE3 and NonO-TFE3 all bind to MET promoter. However, ASPL-TFE3 induces a much stronger up-regulation of downstream MET receptor tyrosine kinase than PSF-TFE3 or NonO-TFE3 does. Evidence shows that the chimeric PRCC-TFE3 and NonO-TFE3 are more potent as transcription factors than wild type TFE3, while PSF-TFE3 and CLTC-TFE3 interfere with cell cycle control.<sup>36</sup>

Table 1. Translocations Identified in Xp11.2 RCC			
Fusion partner	Chromosomal rearrangement	Gene fusion product	Frequency <sup>1</sup>
PRCC <sup>24</sup>	t(X;1)(p11.2;p23)	PRCC-TFE3	70%
PSF <sup>25</sup>	t(X;1)(p11.2;p23)	PSF-TFE3	30%
ASPL <sup>26,27</sup>	t(X;1)(p11.2;p23)	ASPL-TFE3	Rare
NonO <sup>28</sup>	t(X;1)(p11.2;p23)	NonO-TFE3	Rare
CLTC <sup>29,30</sup>	t(X;1)(p11.2;p23)	CLTC-TFE3	Rare
Unknown <sup>31</sup>	t(X;1)(p11.2;p23)	Unknown-TFE3	Rare
Unknown <sup>32</sup>	t(X;1)(p11.2;p23)	Unknown-TFE3	Rare
Unknown <sup>33</sup>	t(X;1)(p11.2;p23)	Unknown-TFE3	Rare


[Click icon to read this table](#)

**Table 1.** Translocations Identified in Xp11.2 RCC.

## HISTOPATHOLOGY


Grossly, Xp11.2 RCC is indistinguishable from conventional RCC.<sup>17,28</sup> The tumors are well circumscribed, but unencapsulated with tan-yellow, soft cut surfaces. Tumor size varies from 2.1 to 21 cm with mean size of 6.8 cm.<sup>15</sup> Areas with necrosis, hemorrhage, calcification and cystic changes may be present. Histologically, Xp11.2 RCC has nested or tubular to papillary growth patterns. Tumor cells have distinctive clear to eosinophilic, voluminous, granular cytoplasm and prominent cell borders. The nuclei are vesicular and have prominent nucleoli. Psammomatous calcification is also a common feature (**Figure 1**).<sup>2,28,37</sup> The morphology of Xp11.2 RCC with different gene fusion partners may vary slightly. For example, ASPL-*TFE3* RCC is associated with exuberant psammomatous calcification and such abundant cytoplasm that it was called “voluminous cell variant” of pediatric RCC before the translocation was identified. In contrast, the cytoplasm is less abundant and the psammoma bodies are fewer in PRCC-*TFE3* RCC. The histology variation might be explained by the heterogeneity of fusion partners of *TFE3*. Whether subtle morphologic differences exist in other Xp11.2 RCCs, such as PSF-*TFE3*, NonO-*TFE3*, CLTC-*TFE3* is currently not clear.<sup>11</sup> Because histologically Xp11.2 RCCs have clear cells with distinctive cell borders and grow in nested and/or tubular-papillary patterns, they may resemble conventional clear cell RCC (CCRCC), papillary RCC (PRCC), and clear cell papillary RCC (CCPRCC), a recently recognized new entity. In addition to the unique morphological features of each entities mentioned above, immunohistochemical markers are helpful in the differential diagnoses. Unlike other types of RCC, Xp11.2 RCC is negative or only focally positive for cytokeratins and epithelial membrane antigen (EMA). Vimentin is usually negative but may be weakly and focally positive. Nuclear staining of *TFE3* is highly sensitive and specific for Xp11.2 RCC (**Figure 2**).<sup>38</sup> CD10 and alpha-methylacyl-coenzyme A racemase (AMACR) are also commonly positive in Xp11.2 RCC, but not specific.<sup>39</sup> In addition, recent studies by Argani et al.<sup>14</sup> showed that Xp11.2 RCCs rarely express c-kit; but they show high levels of phosphorylated S6, which is a mTOR pathway activation marker. In difficult cases with equivocal histology and immunohistochemical stainings, break-apart fluorescence in situ hybridization (FISH) assay and reverse transcriptase-polymerase chain reaction (RT-PCR) are useful confirmatory tests in the diagnosis of Xp11.2 RCC.<sup>17,40-41</sup> The immunohistochemical staining pattern for Xp11.2 RCC and other RCCs is summarized in **Table 2**.

Go To: | [Top](#) | [Abstract](#) | [Text](#) | [Author Information](#) | [References](#) |



[Click icon to view larger version](#)

**Figure 1.** Characteristic microscopic features of Xp11.2 RCC. **A.** Papillary growth pattern with mixed clear/eosinophilic cells (H&E, X100); **B.** Nested growth pattern with mixed clear/eosinophilic cells (H&E, X100); **C.** Psammoma bodies are common (H&E, X100); **D.** Tumor cells have voluminous cytoplasm and distinct cell borders. Nuclei are vesicular and have prominent nucleoli (H&E, X400).



[Click icon to view larger version](#)

**Figure 2.** Positive nuclear staining for *TFE3* in Xp11.2 RCC (X400). The nuclei of tumor cells are positive for *TFE3* using an antibody against the C-terminal portion of the *TFE3* in tubular/nested (**A**) and papillary (**B**) areas.

## PROGNOSIS AND TREATMENT

Reports regarding the prognosis of Xp11.2 RCC in children and young adults are controversial. Initially, it was believed that the biological behavior of Xp11.2 RCC is indolent. In a study conducted by Ramphal et al.,<sup>25</sup> 13 pediatric patients diagnosed with stage I to IV Xp11.2 RCC underwent nephrectomy and resection of metastases with negative surgical margins. All but one patients were alive and tumor-free at last follow-up with an overall survival rates of  $92\% \pm 7.4\%$  at 5 years. However, recently, both prospective and retrospective studies have shown that Xp11.2 RCC is associated with significantly decreased disease-free survival and overall survival in this group of patients.<sup>42-43</sup> In adults, Xp11.2 RCC has a more aggressive clinical course with advanced stage at diagnosis, development of hematogenous metastases and rapid relapse.<sup>15,17,44-45</sup> The cancer-specific survival rate is significantly decreased in patients with Xp11.2 RCC than those with other types of RCCs.<sup>45</sup> The grim clinical outcome associated with Xp11.2 RCC warrants early detection, accurate diagnosis and close follow-up. The current management of Xp11.2 RCC is similar to conventional RCC. For localized Xp11.2 RCC including patients with positive regional lymph nodes, surgery is the treatment of choice.<sup>20,46</sup> For patients with hematogenous metastases, the current options are immunotherapy using cytokines, such as interleukin 2 (IL-2) and interferon alpha (IFN $\alpha$ ) and multi-kinase inhibitors. Clinical studies have demonstrated certain efficacy of multi-kinase inhibitors, for example, sunitinib, sorafenib and mTOR/MET kinase inhibitor, in treating rapidly progressive metastatic Xp11.2 RCC in adult patients.<sup>47-53</sup> The optimal treatment approach for Xp11.2 RCC remains to be determined.

Table 2. The Immunostain Profiles of Xp11.2 RCC and its Close Mimickers.

	Xp11.2 RCC	CCRCC	PRCC	CCPRCC
TFE3	+	-	-	-
Cathepsin K	+	-	-	-
Cyt	-	+	+	-
Vimentin	-	+	+	- or focal +
AMACR	+	-	-	-
CD10	+	+	+	- or focal +
CA9	-	+	+	+

**Table 2.** The Immunostain Profiles of Xp11.2 RCC and its Close Mimickers.

[Click icon to read this table](#)

## SUMMARY

Xp11.2 RCC is a rare tumor affecting both pediatric and adult populations and is relatively common in children and adolescents. The clinical presentation is nonspecific and patients typically have a poor prognosis. In children, prior chemotherapy may predispose to developing Xp11.2 RCC. Eight TFE3 fusion partners have been reported; however, the identities of three of them are unknown. The chimeric TFE3 fusion proteins contribute to tumorigenesis by dysregulating gene transcription and cell cycle. The characteristic histological pattern of Xp11.2 RCC is clear cells with voluminous cytoplasm arranged in nested and/or tubular-papillary architecture. Its immunohistochemical staining profile is unique, showing positive nuclear staining for TFE3 and Cathepsin-K and negative or focally weakly positive for cytokeratins. Modern molecular techniques of FISH and RT-PCR can confirm the diagnosis of Xp11.2 RCC. Surgery remains to be the treatment of choice for organ confined or cases with limited local metastasis. Newer therapeutic approach targeting the aberrant MiTF/TFE transcriptional pathway, such as mTOR/MET inhibitors, may provide alternative treatment in the future.

Go To: | [Top](#) | [Abstract](#) | [Text](#) | [Author Information](#) | [References](#) |

**Yunguang Liu, MD, PhD;<sup>1</sup> Bo Xu, MD, PhD;<sup>2</sup> Frank Chen, MD, PhD<sup>1\*</sup>**

<sup>1</sup> Department of Pathology, Buffalo General Hospital, State University of New York, Buffalo, NY

<sup>2</sup> Department of Pathology and Laboratory Medicine, Roswell Park Cancer Institute, Buffalo, NY

**\*Corresponding Author:** Department of Pathology, Buffalo General Hospital, 100 High Street, State University of New York, Buffalo, NY 14203. (Email: dr.frankchen@yahoo.com).

## CONFLICT OF INTEREST

The authors have no conflicts of interest to disclose.

Go To: | [Top](#) | [Abstract](#) | [Text](#) | [Author Information](#) | [References](#) |

## REFERENCES

- < 1. de Jong B, Molenaar IM, Leeuw JA, Idenberg VJ, Oosterhuis JW. Cytogenetics of a renal adenocarcinoma in a 2-year-old child. *Cancer Genet Cytogenet.* 1986;21(2):165-169.
- < 2. Argani P, Ladanyi M. Renal carcinomas associated with Xp11.2 translocations/TFE3 gene fusions. In: Eble JN, Sauter G, Epstein JI, et al, eds. *Pathology and Genetics of Tumours of the Urinary System and Male Genital Organs (World Health Organization Classification of Tumours)*. Lyon, France: IARC Press; 2004:37-38.
- < 3. Fisher DE, Carr CS, Parent LA, Sharp PA. TFEB has DNA-binding and oligomerization properties of a unique helix-loop-helix/leucine-zipper family. *Genes Dev.* 1991;5(12A):2342-2352.
- < 4. Hemesath TJ, Steingrimsson E, McGill G, et al. Microphthalmia, a critical factor in melanocyte development, defines a discrete transcription factor family. *Genes Dev.* 1994;8(22):2770-2780.
- < 5. Kuiper RP, Schepens M, Thijssen J, Schoenmakers EF, van Kessel AG. Regulation of the MiTF/TFE bHLH-LZ transcription factors through restricted spatial expression and alternative splicing of functional domains. *Nucleic Acids Res.* 2004;32(8):2315-2322.
- < 6. Zhao GQ, Zhao Q, Zhou X, Mattei MG, de Crombrughe B. TFEC, a basic helix-loop-helix protein, forms heterodimers with TFE3 and inhibits TFE3-dependent transcription activation. *Mol Cell Biol.* 1993;13(8):4505-4512.
- < 7. Davis IJ, Fisher DE. MiT transcription factor associated malignancies in man. *Cell Cycle.* 2007;6(14):1724-1729.
- < 8. Argani P, Lal P, Hutchinson B, Lui MY, Reuter VE, Ladanyi M. Aberrant nuclear immunoreactivity for TFE3 in neoplasms with TFE3 gene fusions: a sensitive and specific immunohistochemical assay. *Am J Surg Pathol.* 2003;27(6):750-761.
- < 9. Ries L, Smith M, Gurney J, et al, eds. *Cancer Incidence and Survival among Children and Adolescents: United States SEER Program 1975-1995* Bethesda, MD: National Cancer Institute;1999.
- < 10. Bruder E, Passera O, Harms D, et al. Morphologic and molecular characterization of renal cell carcinoma in children and young adults. *Am J Surg Pathol.* 2004;28(9):1117-1132.
- < 11. Argani P, Ladanyi M. Translocation carcinomas of the kidney. *Clin Lab Med.* 2005;25(2):363-378.
- < 12. Macher-Goeppinger S, Wilfried R, Wagener N, et al. Molecular heterogeneity of TFE3 activation in renal cell carcinomas. *Mod Pathol.* 2011;169. [Epub ahead of print].
- < 13. Tomlinson GE, Nisen PD, Timmons CF, Schneider NR. Cytogenetics of a renal cell carcinoma in a 17-month-old child. Evidence for Xp11.2 as a recurring breakpoint. *Cancer Genet Cytogenet.* 1991;57(1):11-17.
- < 14. Argani P, Hicks J, De Marzo AM, et al. Xp11 translocation renal cell carcinoma (RCC): extended immunohistochemical profile emphasizing novel RCC markers. *Am J Surg Pathol.* 2010;34(9):1295-1303.
- < 15. Argani P, Olgac S, Tickoo SK, et al. Xp11 translocation renal cell carcinoma in adults: expanded clinical, pathologic, and genetic spectrum. *Am J Surg Pathol.* 2007;31(8):1149-1160.
- < 16. Haudebourg J, Hoch B, Fabas T, et al. A novel case of t(X;1)(p11.2;p34) in a renal cell carcinoma with TFE3 rearrangement and favorable outcome in a 57-year-old patient. *Cancer Genet Cytogenet.* 2010;200(2):75-78.
- < 17. Hung CC, Pan CC, Lin CC, Lin AT, Chen KK, Chang YH. XP11.2 Translocation renal cell carcinoma: Clinical experience of Taipei Veterans General Hospital. *J Chin Med Assoc.* 2011;74(11):500-504.
- < 18. Argani P, Ladanyi M. Distinctive neoplasms characterised by specific chromosomal translocations comprise a significant proportion of paediatric renal cell carcinomas. *Pathology.* 2003;35(6):492-498.
- < 19. Prasad SR, Humphrey PA, Catena JR, et al. Common and uncommon histologic subtypes of renal cell carcinoma: imaging spectrum with pathologic correlation. *Radiographics.* 2006;26(6):1795-1806; discussion 1806-1710.
- < 20. Horie K, Kikuchi M, Miwa K, et al. A case of xp11.2 translocation renal cell carcinoma. *Hinyokika Kiyo.* 2011;57(3):129-133.
- < 21. Kato H, Kanematsu M, Yokoi S, et al. Renal cell carcinoma associated with Xp11.2 translocation/TFE3 gene fusion: radiological findings mimicking papillary subtype. *J Magn Reson Imaging.* 2011;33(1):217-220.
- < 22. Komai Y, Fujiwara M, Fujii Y, et al. Adult Xp11 translocation renal cell carcinoma diagnosed by cytogenetics and immunohistochemistry. *Clin Cancer Res.* 2009;15(4):1170-1176.
- < 23. Argani P, Lae M, Ballard ET, et al. Translocation carcinomas of the kidney after chemotherapy in childhood. *J Clin Oncol.* 2006;24(10):1529-1534.
- < 24. Brassesco MS, Valera ET, Bonilha TA, et al. Secondary PSF/TFE3-associated renal cell carcinoma in a child treated for genitourinary rhabdomyosarcoma. *Cancer Genet.* 2011;204(2):108-110.
- < 25. Ramphal R, Pappo A, Zielenska M, Grant R, Ngan BY. Pediatric renal cell carcinoma: clinical, pathologic, and molecular abnormalities associated with the members of the mit transcription factor family. *Am J Clin Pathol.* 2006;126(3):349-364.

- < 26. Armah HB, Parwani AV, Surti U, Bastacky SI. Xp11.2 translocation renal cell carcinoma occurring during pregnancy with a novel translocation involving chromosome 19: a case report with review of the literature. *Diagn Pathol.* 2009;4:15.
- < 27. Nouh MA, Kuroda N, Yamashita M, et al. Renal cell carcinoma in patients with end-stage renal disease: relationship between histological type and duration of dialysis. *BJU Int.* 2010;105(5):620-627.
- < 28. Argani P, Antonescu CR, Illei PB, et al. Primary renal neoplasms with the ASPL-TFE3 gene fusion of alveolar soft part sarcoma: a distinctive tumor entity previously included among renal cell carcinomas of children and adolescents. *Am J Pathol.* 2001;159(1):179-192.
- < 29. Argani P, Lui MY, Couturier J, Bouvier R, Fournet JC, Ladanyi M. A novel CLTC-TFE3 gene fusion in pediatric renal adenocarcinoma with t(X;17)(p11.2;q23). *Oncogene.* 2003;22(34):5374-5378.
- < 30. Dijkhuizen T, van den Berg E, Wilbrink M, et al. Distinct Xp11.2 breakpoints in two renal cell carcinomas exhibiting X;autosome translocations. *Genes Chromosomes Cancer.* 1995;14(1):43-50.
- < 31. Murphy WM, Grignon DJ, Perlman EJ, eds. *Tumors of the Kidney, Bladder and Related Urinary Structures 2004 (AFIP Atlas of Tumor Pathology 4th Series).* 4th ed. American Registry of Pathology, publisher; 2004.
- < 32. Ladanyi M, Lui MY, Antonescu CR, et al. The der(17)t(X;17)(p11;q25) of human alveolar soft part sarcoma fuses the TFE3 transcription factor gene to ASPL, a novel gene at 17q25. *Oncogene.* 2001;20(1):48-57.
- < 33. Martignoni G, Pea M, Gobbo S, et al. Cathepsin-K immunoreactivity distinguishes MiTF/TFE family renal translocation carcinomas from other renal carcinomas. *Mod Pathol.* 2009;22(8):1016-1022.
- < 34. Argani P. The evolving story of renal translocation carcinomas. *Am J Clin Pathol.* 2006;126(3):332-334.
- < 35. Tsuda M, Davis IJ, Argani P, et al. TFE3 fusions activate MET signaling by transcriptional up-regulation, defining another class of tumors as candidates for therapeutic MET inhibition. *Cancer Res.* 2007;67(3):919-929.
- < 36. Medendorp K, van Groningen JJ, Schepens M, et al. Molecular mechanisms underlying the MiT translocation subgroup of renal cell carcinomas. *Cytogenet Genome Res.* 2007;118(2-4):157-165.
- < 37. Argani P, Antonescu CR, Couturier J, et al. PRCC-TFE3 renal carcinomas: morphologic, immunohistochemical, ultrastructural, and molecular analysis of an entity associated with the t(X;1)(p11.2;q21). *Am J Surg Pathol.* 2002;26(12):1553-1566.
- < 38. Ross H, Argani P. Xp11 translocation renal cell carcinoma. *Pathology.* 2010;42(4):369-373.
- < 39. Camparo P, Vasiliu V, Molinie V, et al. Renal translocation carcinomas: clinicopathologic, immunohistochemical, and gene expression profiling analysis of 31 cases with a review of the literature. *Am J Surg Pathol.* May 2008;32(5):656-670.
- < 40. Kim SH, Choi Y, Jeong HY, Lee K, Chae JY, Moon KC. Usefulness of a break-apart FISH assay in the diagnosis of Xp11.2 translocation renal cell carcinoma. *Virchows Arch.* 2011;459(3):299-306.
- < 41. Zhong M, De Angelo P, Osborne L, et al. Dual-color, break-apart FISH assay on paraffin-embedded tissues as an adjunct to diagnosis of Xp11 translocation renal cell carcinoma and alveolar soft part sarcoma. *Am J Surg Pathol.* 2010;34(6):757-766.
- < 42. Rao Q, Chen JY, Wang JD, et al. Renal cell carcinoma in children and young adults: clinicopathological, immunohistochemical, and VHL gene analysis of 46 cases with follow-up. *Int J Surg Pathol.* 2011;19(2):170-179.
- < 43. Rao Q, Guan B, Zhou X. Xp11.2 Translocation Renal Cell Carcinomas Have a Poorer Prognosis Than Non-Xp11.2 Translocation Carcinomas in Children and Young Adults: A Meta-analysis. *Int J Surg Pathol.* 2010(18):458-464.
- < 44. Meyer PN, Clark JJ, Flanigan RC, Picken MM. Xp11.2 translocation renal cell carcinoma with very aggressive course in five adults. *Am J Clin Pathol.* 2007;128(1):70-79.
- < 45. Mir MC, Trilla E, de Torres IM, et al. Altered transcription factor E3 expression in unclassified adult renal cell carcinoma indicates adverse pathological features and poor outcome. *BJU Int.* 2011;108(2 Pt 2):E71-76.
- < 46. Aoyagi T, Shinohara N, Kubota-Chikai K, Kuroda N, Nonomura K. Long-term survival in a patient with node-positive adult-onset Xp11.2 translocation renal cell carcinoma. *Urol Int.* 2011;86(4):487-490.
- < 47. Choueiri TK, Lim ZD, Hirsch MS, et al. Vascular endothelial growth factor-targeted therapy for the treatment of adult metastatic Xp11.2 translocation renal cell carcinoma. *Cancer.* 2010;116(22):5219-5225.
- < 48. Hou MM, Hsieh JJ, Chang NJ, et al. Response to sorafenib in a patient with metastatic xp11 translocation renal cell carcinoma. *Clin Drug Investig.* 2010;30(11):799-804.
- < 49. Liu YC, Chang PM, Liu CY, Yang CY, Chen MH, Pan CC. Sunitinib-induced nephrotic syndrome in association with drug response in a patient with Xp11.2 translocation renal cell carcinoma. *Jpn J Clin Oncol.* 2011;41(11):1277-1281.
- < 50. Numakura K, Tsuchiya N, Yuasa T, et al. A case study of metastatic Xp11.2 translocation renal cell carcinoma effectively treated with sunitinib. *Int J Clin Oncol.* 2011;16(5):577-580.

- < 51. Parikh J, Coleman T, Messias N, Brown J. Temsirolimus in the treatment of renal cell carcinoma associated with Xp11.2 translocation/TFE gene fusion proteins: a case report and review of literature. *Rare Tumors*. 2009;1(2):e53.
- < 52. Pwint TP, Macaulay V, Roberts IS, Sullivan M, Protheroe A. An adult Xp11.2 translocation renal carcinoma showing response to treatment with sunitinib. *Urol Oncol*. 2011;29(6):821-824.
- < 53. Wong MK, Jarkowski A. Response to sorafenib after sunitinib-induced acute heart failure in a patient with metastatic renal cell carcinoma: case report and review of the literature. *Pharmacotherapy*. 2009;29(4):473-478.
- .....

Go To: | [Top](#) | [Abstract](#) | [Text](#) | [Author Information](#) | [References](#) |

---

Comments are closed.

Copyright © 2014 [NAJMS: The North American Journal of Medicine and Science](#). All Rights Reserved.

Magazine Basic theme designed by [Themes by bavotasan.com](#).

Powered by [WordPress](#).